

Qualitative Analysis of the Nonlinear Double Degenerate Parabolic Equation of  
Turbulent Filtration with Absorption

by

Adam Prinkey

Bachelor of Science  
Biochemistry (with a focus on Biology)  
Florida Institute of Technology  
2013

Bachelor of Science  
Mathematical Sciences  
Florida Institute of Technology  
2013

A dissertation  
submitted to Florida Institute of Technology  
in partial fulfillment of the requirements  
for the degree of

Doctorate of Philosophy  
in  
Applied Mathematics

Melbourne, Florida  
May, 2019

© Copyright 2019 Adam Prinkey

All Rights Reserved

---

The author grants permission to make single copies.

We the undersigned committee  
hereby approve the attached dissertation

Qualitative Analysis of the Nonlinear Double Degenerate Parabolic Equation of  
Turbulent Filtration with Absorption by

Adam Prinkey

---

Ugur G. Abdulla, Ph.D., Dr.Sci., Dr.rer.nat.habil.  
Professor  
Department of Mathematical Sciences  
Committee Chair

---

David Carroll, Ph.D.  
Associate Professor  
Department of Biomedical and Chemical Engi-  
neering and Sciences  
Outside Committee Member

---

Jian Du, Ph.D.  
Associate Professor  
Department of Mathematical Sciences  
Committee Member

---

Tariel Kiguradze, Ph.D.  
Associate Professor  
Department of Mathematical Sciences  
Committee Member

---

Munevver Subasi, Ph.D.  
Associate Professor and Department Head  
Department of Mathematical Sciences

## ABSTRACT

Title:

Qualitative Analysis of the Nonlinear Double Degenerate Parabolic Equation of  
Turbulent Filtration with Absorption

Author:

Adam Prinkey

Major Advisor:

Ugur G. Abdulla, Ph.D., Dr.Sci., Dr.rer.nat.habil.

The goal of the dissertation is to pursue qualitative analysis of the mathematical model of turbulent polytropic filtration of a gas in a porous media with reaction or absorption described by the second order nonlinear double degenerate parabolic equation

$$\frac{\partial u}{\partial t} - \frac{\partial}{\partial x} F \left[ \frac{\partial u^m}{\partial x} \right] + Q(u) = 0, \quad (1)$$

where

$$F(y) = |y|^{p-1}y, Q(u) = bu^\beta, m, p, \beta > 0, b \in \mathbb{R}.$$

In the absence of the reaction term there is a finite speed of propagation with an expanding interface in the case of slow diffusion ( $mp > 1$ ), and infinite speed of propagation in the case of fast diffusion ( $0 < mp < 1$ ). In general, qualitative properties of the turbulent filtration is an outcome of the competition between the diffusion and reaction or absorption forces. In the slow diffusion case, the strong domination of the diffusion causes an expanding interface. In the fast diffusion case, the strong domination of the diffusion causes infinite speed of propagation and absence of interfaces, while weak domination of the diffusion causes an expanding interface. Domination of the absorption causes a shrinking interface. If diffusion and absorption are in balance then initial density pro-

file dictates the direction of the interface movement. When the interface exists, explicit asymptotic formulas and estimates for the interface function and the local solution near the interfaces with accuracy up to precise constants are proved. Explicit asymptotic formulas for the local solution at infinity are proved in all cases where interface does not exist. The results of the dissertation can be applied to problems in the oil and gas industry to pursue the estimation and control of the time evolution of the size of oil and gas resources.

# Table of Contents

<b>Abstract</b>	<b>iii</b>
<b>List of Figures</b>	<b>viii</b>
<b>List of Notations</b>	<b>ix</b>
<b>Acknowledgments</b>	<b>xii</b>
<b>Dedication</b>	<b>xiii</b>
<b>1 Introduction</b>	<b>1</b>
1.1 Physical Derivation of the Equation . . . . .	1
1.1.1 The Barenblatt Solution . . . . .	3
1.2 Historical Review and the Open Problem . . . . .	3
1.2.1 The Open Problem . . . . .	4
<b>2 Evolution of Interfaces for the Nonlinear Double Degenerate Parabolic Equation with Slow Diffusion</b>	<b>7</b>
2.1 Description of the Main Results . . . . .	8
2.2 Further Details of the Main Results . . . . .	10
2.3 Preliminary Results . . . . .	15

2.3.1	Asymptotic Properties of Solutions With $mp > 1$ . . . . .	17
2.3.1.1	Diffusion Dominates Reaction . . . . .	17
2.3.1.2	Diffusion and Reaction are in Balance . . . . .	24
2.3.1.3	Reaction Dominates Diffusion . . . . .	27
2.4	Proofs of the Main Results . . . . .	29
2.4.1	Proof of Theorem 2.1.1 . . . . .	29
2.4.2	Proof of Theorem 2.1.2 . . . . .	30
2.4.3	Proof of Theorem 2.1.3 . . . . .	36

<b>3</b>	<b>Traveling-Wave Solutions to the Nonlinear Double Degenerate Reaction-Diffusion Equation</b>	<b>40</b>
3.1	Introduction and the Main Result . . . . .	40
3.1.1	Preliminary Results: Traveling-Wave Solutions and Phase-Space Analysis . . . . .	42
3.2	Proof of the Main Result . . . . .	55
<b>4</b>	<b>Evolution of Interfaces for the Nonlinear Double Degenerate Parabolic Equation with Fast Diffusion</b>	<b>61</b>
4.1	Main Results . . . . .	61
4.2	Additional Details of the Results . . . . .	65
4.3	Asymptotic Properties of Solutions With $0 < mp < 1$ . . . . .	69
4.3.0.1	Diffusion Weakly Dominates Reaction . . . . .	69
4.3.0.2	Diffusion and Reaction are in Balance . . . . .	71
4.3.0.3	Reaction Dominates Diffusion . . . . .	73
4.4	Proofs of the Main Results . . . . .	73
<b>5</b>	<b>Conclusions</b>	<b>78</b>
	<b>References</b>	<b>82</b>
<b>6</b>	<b>Appendix</b>	<b>95</b>
6.1	. . . . .	95
6.2	. . . . .	97



# List of Figures

2.1	$(\alpha, \beta)$ parameter space diagram for the interface development for the CP (1.7), (1.9) with $mp > 1$ , which is presented in [20]. . . . .	8
3.1	Trajectories $\Upsilon(\Theta)$ and $v(t)$ . . . . .	46
4.1	$(\alpha, \beta)$ parameter space diagram for the interface development for the CP (1.7), (1.9) with $0 < mp < 1$ , which is presented in [24]. . . . .	62

# List of Notations

- $(\cdot)_+ := \max\{\cdot, 0\}$ .
- $C^k(\Omega)$  is the space of  $k$ -times differentiable functions in  $\Omega$  with finite norm

$$\|f\|_{C^k(\Omega)} := \sum_{n=0}^k \sup_{x \in \Omega} |f^{(n)}(x)|.$$

If  $k = 0$ ,  $C^0(\Omega) \equiv C(\Omega)$  denotes the space of continuous functions on  $\Omega$ .

- $C_0^k(\Omega)$  is the space of  $k$ -times differentiable functions in  $\Omega$  with compact support: the space of  $k$ -times differentiable functions in  $\Omega$  that are identically zero outside of a compact (bounded and closed) set.
- $L^p(\Omega)$  ( $1 \leq p < +\infty$ ) is the space of  $p$ -integrable functions in  $\Omega$  with finite norm

$$\|f\|_{L^p(\Omega)} := \left( \int_{\Omega} |f(x)|^p dx \right)^{1/p}.$$

- $L_{\text{loc}}^p(\Omega)$  is the space of locally  $p$ -integrable functions in  $\Omega$ : the space of  $p$ -integrable functions on compact subsets of  $\Omega$ .
- $L^\infty(\Omega)$  is the space of all essentially bounded and measurable functions on  $\Omega$  with

finite norm

$$\|f\|_{L^\infty(\Omega)} := \operatorname{ess\,sup}_{x \in \Omega} |f(x)| = \inf\{C \geq 0 : |f(x)| \leq C, \text{ for almost every } x \in \Omega\}.$$

- Let  $I \subset \mathbb{R}$  and  $u \in L^p(I)$ . The function  $v \in L^p(I)$  is called the weak derivative of  $u$  if the following integral identity holds for all test functions  $\varphi \in C_0^\infty(I)$

$$\int_I u(z) \varphi'(z) dz = - \int_I v(z) \varphi(z) dz.$$

- $W^{k,p}(\Omega)$  ( $1 \leq p \leq +\infty$ ) is the time-independent Sobolev space of functions in  $L^p(\Omega)$  such that the functions themselves and their weak derivatives up to order  $k$  have a finite  $L^p$  norm, with finite Sobolev norm

$$\|f\|_{W^{k,p}(\Omega)} := \left( \sum_{i=0}^k \|f^{(i)}\|_{L^p(\Omega)}^p \right)^{1/p}.$$

- Time-dependent Sobolev spaces. The following spaces are of importance to the study of time-dependent problems, specifically parabolic problems

$$C(0, T; L_{\text{loc}}^1(\Omega)) := \left\{ u = u(t) : [0, T] \rightarrow L_{\text{loc}}^1(\Omega) \right\},$$

let  $K \subset \Omega$  be compact,

$$\|u\|_{C(0, T; L^1(K))} := \max_{0 \leq t \leq T} \|u\|_{L^1(K)}.$$

We define the time-dependent Sobolev space  $L^p(0, T; W_{\text{loc}}^{1,p}(K))$  as

$$L^p(0, T; W_{\text{loc}}^{1,p}(K)) := \left\{ u = u(t) : [0, T] \rightarrow W_{\text{loc}}^{1,p}(K) \right\},$$

with finite norm

$$\|u\|_{L^p(0, T; W^{1,p}(K))} := \left( \int_0^T \|u\|_{W^{1,p}(K)}^p dt \right)^{1/p} = \left( \int_0^T \int_K (|u(x, t)|^p + |Du(x, t)|^p) dx dt \right)^{1/p}.$$

# Acknowledgements

I would like to thank my advisor, Professor Ugur G. Abdulla, for sharing his deep expertise in the field nonlinear partial differential equations, in general, and for his constant patience with me through many meetings over several years. Without his help, this work would not have been possible. I would also like to acknowledge my wife, Megan, and son, Benjamin, for their love, encouragement, and support, throughout the completion of this work. Finally, I would like to thank my mother, Michele, without whom I would not have been able to complete my undergraduate education, and so, would not have been able to pursue my doctorate at all. I'm at this point because of all of you, and I cannot thank you enough for helping me through this challenging journey.

# Dedication

This dissertation is dedicated to my wife, Megan, my son, Benjamin, my mother, Michele, and my mom-mom, Betsy. I love you all very much.

# Chapter 1

## Introduction

### 1.1 Physical Derivation of the Equation

Consider the physical problem of turbulent filtration of a fluid, specifically a gas, in a porous media. This problem is governed by the following physical laws:

- Polytropic equation of state, relating the fluid pressure to its density

$$\mathcal{P} = c\vartheta^n, \tag{1.1}$$

where  $c$  is a positive constant,  $\vartheta = \vartheta(x, t)$  is a density of the fluid, and  $\mathcal{P}$  is the pressure of the fluid.

- Continuity equation, relating the fluid density to its velocity

$$k\vartheta_t + (\vartheta\mathcal{V})_x = 0, \tag{1.2}$$

where  $\mathcal{V}$  is the velocity of the fluid at the point  $x$  and moment  $t$  and  $k$  is a positive

constant.

- Flux under turbulence

$$\vartheta \mathcal{V} = -Q|\phi_x|^{p-1}\phi_x, \quad (1.3)$$

where  $\phi = \mathcal{P}^{\frac{n+1}{n}}$ ,  $Q$  is a positive physical constant, with  $n \geq 1$  and  $p \geq 1$ .

Combining (1.1)-(1.3), we have

$$k\vartheta_t = Qc^{\frac{p(n+1)}{n}} \left( |(\vartheta^{n+1})_x|^{p-1} (\vartheta^{n+1})_x \right)_x. \quad (1.4)$$

By rescaling  $t$  in (1.4) and setting  $\vartheta = u = u(x, t)$  we obtain the nonlinear double degenerate diffusion equation

$$u_t = \left( |(u^m)_x|^{p-1} (u^m)_x \right)_x,$$

where  $m = n + 1 \geq 2$  under these particular conditions, however, this is relaxed throughout the dissertation: we only assume  $m$  and  $p$  are positive constants.

The case where  $mp > 1$  is called the slow diffusion case and the case  $0 < mp < 1$  is called the fast diffusion case [74]. If  $m = 1$ , then we have the non-Newtonian elastic filtration equation or the parabolic  $p$ -Laplacian equation, while if  $m = p = 1$  we have the linear diffusion equation or heat equation.



### 1.1.1 The Barenblatt Solution

Consider the Barenblatt problem for the nonlinear double degenerate parabolic equation, a standard model problem for nonlinear second order degenerate equations

$$\begin{cases} u_t = (|(u^m)_x|^{p-1}(u^m)_x)_x, & x \in \mathbb{R}, t > 0, \\ u(x, 0) = \delta(x), & \forall x \neq 0, \end{cases}$$

where  $\delta(\cdot)$  is the Dirac measure on  $\mathbb{R}$ .

The solution, called the Barenblatt solution, is given by the formula

$$u_*(x, t) = t^{\frac{-1}{p(m+1)}} \left[ \Gamma - k(m, p) \left( |x| t^{\frac{-1}{p(m+1)}} \right)^{\frac{1+p}{p}} \right]_{+}^{\frac{p}{mp-1}}, \quad (1.5)$$

where  $k(m, p) = \frac{mp-1}{m(1+p)} \left( \frac{1}{p(m+1)} \right)^{\frac{1}{p}}$  and  $\Gamma$  is an integration constant defined by the conservation of energy condition. This solution, (1.5), is called the Barenblatt solution. In the slow diffusion case ( $mp > 1$ ) [42, 43], the solution travels with a finite speed of propagation (the interface function exists). In the fast diffusion case ( $0 < mp < 1$ ), solution of the above problem exhibits an infinite speed of propagation. With  $mp > 1$ , the case of slow diffusion, the solution exhibits a finite speed of propagation, and so, is not a classical.

## 1.2 Historical Review and the Open Problem

The solutions to equation (1) are not classical and are understood in the following, *weak*, sense

**Definition 1.2.1.** For the equation (1), we introduce the notion of the weak solution.

A continuous nonnegative function  $u(x, t)$  defined in  $\mathbb{R} \times [0, T)$  is a weak solution of

(1.7), (1.8) if for any  $T_1 \in (0, T)$  and any bounded interval  $(a, b) \in \mathbb{R}$ ,  $(u^m)_x \in L^{p+1}((a, b) \times (0, T_1))$  and

$$\int_0^{T_1} \int_a^b \left( -u\phi_t + |(u^m)_x|^{p-1}(u^m)_x\phi_x + bu^\beta\phi \right) dxdt = \int_a^b u\phi \Big|_{t=T_1}^{t=0} dx \quad (1.6)$$

for arbitrary  $\phi \in C^1([a, b] \times [0, T_1])$  such that  $\phi \Big|_{x=a} = \phi \Big|_{x=b} = 0$ .

The Barenblatt solution above,  $u_*$ , when  $mp > 1$ , is a weak solution in the sense of the Definition 1.2.1 due to having a finite speed of propagation.

The general theory of nonlinear degenerate parabolic equations is well studied (refer to [6, 11, 2, 5, 3, 78, 97, 85, 40, 73, 72, 87, 94, 52, 61, 55, 51, 93]). For existence of solutions to general doubly nonlinear parabolic problems we refer to [45, 47, 83]. For advances such as the analysis of fine properties of solutions and the development of Harnack-type inequalities for general doubly nonlinear parabolic problems, refer to [70, 95, 89, 50, 82, 101, 58, 49, 76, 79, 99, 96, 69, 91, 77, 59, 98, 60, 48, 39, 88, 56, 46, 84, 37, 90, 36, 86, 81, 102, 38, 41, 100, 83].

Problems of existence, uniqueness of solutions of general parabolic initial-boundary value problems, the formulation of comparison results, analysis of the regularity properties of weak solutions, and the optimal control of free boundary problems are considered in [30], [18], [19], [42].

### 1.2.1 The Open Problem

We consider the Cauchy problem (CP) for the nonlinear degenerate parabolic equation

$$Lu \equiv u_t - (|(u^m)_x|^{p-1}(u^m)_x)_x + bu^\beta = 0, \quad x \in \mathbb{R}, \quad 0 < t < T, \quad (1.7)$$

with nonnegative and continuous initial function

$$u(x, 0) = u_0(x), \quad x \in \mathbb{R}, \quad (1.8)$$

where  $u = u(x, t)$ ;  $m, p, \beta > 0$ ,  $b \in \mathbb{R}$ , and  $T \leq +\infty$ , with the conditions that if  $b > 0$  then  $\beta < 1$  while if  $b \leq 0$  then  $\beta \geq 1$ . Notably, equation (1.7) models turbulent polytropic filtration of a gas (or general fluid) in a porous medium [42, 57, 78]. We define the interface function as

$$\eta(t) := \sup\{x : u(x, t) > 0\},$$

with  $\eta(0) = 0$ . We want to analyze the short-time behavior of the interface function  $\eta(t)$ , and local solution near the interface function. In general, we shall assume

$$u_0(x) \sim C(-x)_+^\alpha, \text{ as } x \rightarrow 0^-, \text{ for some } C > 0, \alpha > 0. \quad (1.9)$$

In some special cases, in global cases in particular, we will consider

$$u_0(x) = C(-x)_+^\alpha, \quad x \in \mathbb{R}. \quad (1.10)$$

This problem is well understood in the case of (1.7) with  $p = 1$

$$u_t - (u^m)_{xx} + bu^\beta = 0 \quad x \in \mathbb{R}, 0 < t < T. \quad (1.11)$$

Solutions and regularity analysis for the Cauchy problem (1.11), (1.8) is well studied [1, 4, 7, 8, 9, 10, 12, 13, 14, 15, 16, 17, 25, 27, 28, 29, 31, 32, 34, 35, 62, 63]. If  $u_0 \in C(\mathbb{R}) \cap L^\infty(\mathbb{R})$  and nonnegative, then the existence, uniqueness and comparison theorems for the weak solution of the CP (1.7), (1.8) was proved in [57] for the case

$b = 0$ , and in [92] for  $b > 0$ . In [57] it is proved that the weak solution of (1.7),  $b = 0$ , is locally Hölder continuous. Local Hölder continuity of the locally bounded weak solutions of the general second order multidimensional nonlinear degenerate parabolic equations with double degenerate diffusion term is proved in [68, 67].

The interface problem for the CP (1.11), (1.9) was presented in [23] for the slow diffusion case ( $m > 1$ ) and in [33] for the fast diffusion case ( $0 < m < 1$ ). A similar classification for the reaction-diffusion equation with  $p$ -Laplacian type diffusion ( $m = 1$  in (1.7)) is presented in the recent paper [22] for the case of slow diffusion case ( $p > 1$ ) and in [21] for the fast diffusion case ( $0 < p < 1$ ).

The overall goal of this dissertation is to apply and extend the analytical methods developed in [23, 33, 6] to prove accurate asymptotic estimations for the interface function and local solution near the interface function for the Cauchy problem (1.7), (1.8) in the case of both slow diffusion ( $mp > 1$ ) and fast diffusion ( $0 < mp < 1$ ).

## **Chapter 2**

# **Evolution of Interfaces for the Nonlinear Double Degenerate Parabolic Equation with Slow Diffusion**

In this chapter we present full classification of the evolution of interfaces and local structure of solution near the interfaces of the problem (1.7), (1.9) in the slow diffusion case ( $mp > 1$ ). The results of Chapter 2 are published in [20].

## 2.1 Description of the Main Results




Figure 2.1:  $(\alpha, \beta)$  parameter space diagram for the interface development for the CP (1.7), (1.9) with  $mp > 1$ , which is presented in [20].

Throughout this section we assume that  $u$  is a unique weak solution of the CP (1.7)-(1.9). There are four different subcases; the main results are outlined below in Theorems 2.1.1, 2.1.2, 2.1.3 and 2.1.4 corresponding directly to the cases I, II, III and IVa-IVd in Figure 2.1.

**Theorem 2.1.1.** *If  $b > 0$  and  $0 < \alpha < \frac{1+p}{mp-\min\{1,\beta\}}$ , then the interface initially expands and*

$$\eta(t) \sim \xi_* t^{1/(1+p-\alpha(mp-1))}, \text{ as } t \rightarrow 0^+, \quad (2.1)$$

where

$$\xi_* = C^{\frac{mp-1}{1+p-\alpha(mp-1)}} \xi'_*, \quad (2.2)$$

and  $\xi'_*$  is a positive number depending only on  $m$ ,  $p$ , and  $\alpha$ . For arbitrary  $\rho < \xi'_*$ , there exists a positive number  $f(\rho)$  depending on  $C, m, p$ , and  $\alpha$  such that

$$u(x, t) \sim t^{\alpha/(1+p-\alpha(mp-1))} f(\rho), \text{ as } t \rightarrow 0^+, \quad (2.3)$$

along the curve  $x = \xi_\rho(t) = \rho t^{1/(1+p-\alpha(mp-1))}$ .

**Theorem 2.1.2.** *Let  $b > 0$ ,  $0 < \beta < 1$ ,  $\alpha = (1+p)/(mp-\beta)$ , and*

$$C_* = \left[ \frac{b(mp-\beta)^{1+p}}{(m(1+p))^p p(m+\beta)} \right]^{\frac{1}{mp-\beta}}.$$

*Then interface expands or shrinks according as  $C > C_*$  or  $C < C_*$  and*

$$\eta(t) \sim \zeta_* t^{\frac{mp-\beta}{(1+p)(1-\beta)}}, \text{ as } t \rightarrow 0^+, \quad (2.4)$$

*where  $\zeta_* \leq 0$  if  $C \leq C_*$ , and for arbitrary  $\rho < \zeta_*$  there exists  $f_1(\rho) > 0$  such that*

$$u(x, t) \sim t^{1/(1-\beta)} f_1(\rho), \text{ for } x = \rho t^{\frac{mp-\beta}{(1+p)(1-\beta)}}, \text{ as } t \rightarrow 0^+. \quad (2.5)$$

**Theorem 2.1.3.** *If  $b > 0$ ,  $0 < \beta < 1$ , and  $\alpha > (1+p)/(mp-\beta)$ , then the interface shrinks and*

$$\eta(t) \sim -\ell_* t^{1/\alpha(1-\beta)}, \text{ as } t \rightarrow 0^+, \quad (2.6)$$

*where  $\ell_* = C^{-1/\alpha}(b(1-\beta))^{1/\alpha(1-\beta)}$ . For arbitrary  $\ell > \ell_*$  we have*

$$u(x, t) \sim \left[ C^{1-\beta} (-x)_+^{\alpha(1-\beta)} - b(1-\beta)t \right]^{1/(1-\beta)}, \text{ as } t \rightarrow 0^+, \quad (2.7)$$

*along the curve  $x = \eta_\ell(t) = -\ell t^{1/\alpha(1-\beta)}$ .*

**Theorem 2.1.4.** *If  $b \in \mathbb{R}$ ,  $\beta \geq 1$ , and  $\alpha \geq (1+p)/(mp-1)$ , then interface initially remains stationary.*

## 2.2 Further Details of the Main Results

In this section we outline some essential details of the main results described in Theorems 2.1.1-2.1.4 of Section 2.1. In particular, we describe global estimations for minimal solutions of the CP (1.7), (1.10).

*Further details of Theorem 2.1.1.*  $f$  is a shape function of the self-similar solution to the problem (1.7), (1.10) with  $b = 0$

$$u(x, t) = t^{\frac{\alpha}{1+p-\alpha(mp-1)}} f(\xi), \quad \xi = xt^{-\frac{1}{1+p-\alpha(mp-1)}}. \quad (2.8)$$

In fact,  $f$  is a solution of the following nonlinear ODE problem in  $\mathbb{R}$

$$(|(f^m(\xi))'|^{p-1}(f^m(\xi))')' + (1+p-\alpha(mp-1))^{-1}(\xi f'(\xi) - \alpha f(\xi)) = 0, \quad (2.9)$$

$$f(\xi) \sim C(-\xi)^\alpha, \text{ as } \xi \downarrow -\infty, \quad f(+\infty) = 0. \quad (2.10)$$

Moreover, there exists  $\xi_* > 0$  such that:  $f(\xi) \equiv 0$  for  $\xi \geq \xi_*$ ;  $f(\xi) > 0$  for  $\xi < \xi_*$ . Dependence on  $C$  is given by the following relation

$$f(\rho) = C^{1+p/(1+p-\alpha(mp-1))} f_0\left(C^{(mp-1)/(\alpha(mp-1)-(1+p))}\rho\right), \quad (2.11a)$$

$$f_0(\rho) = w(\rho, 1), \quad \xi_*' = \sup\{\rho : f_0(\rho) > 0\} > 0, \quad (2.11b)$$

where  $w$  is a minimal solution of the CP (1.7), (1.10) with  $b = 0$ ,  $C = 1$ . Lower and upper estimations for  $f$  are given in (2.30). We also have that

$$\xi_*' = A_0^{\frac{mp-1}{1+p}} \left[ \frac{(mp)^p(1+p-\alpha(mp-1))}{(mp-1)^p} \right]^{\frac{1}{1+p}} \xi_*'', \quad (2.12)$$



where  $A_0 = w(0, 1)$  and  $\xi''$  is some number belonging to the segment  $[\xi_1, \xi_2]$  (see 6.1). In the particular case  $\alpha = p(mp - 1)^{-1}$  and  $mp > 1 + p - p(\min\{1, \beta\})$ , the explicit solution of (1.7), (1.10) with  $b = 0$  is given by (2.28) and

$$\xi_1 = \xi_2 = 1, \xi_*' = (mp)^p (mp - 1)^{-p}, f_0(x) = (\xi_*' - x)_+^{p/(mp-1)} \quad (2.13)$$

*Further details of Theorem 2.1.2.* If  $p(m + \beta) = 1 + p$ , the solution to (1.7), (1.10) is

$$u(x, t) = C(\zeta_* t - x)_+^{\frac{1+p}{mp-\beta}}, \zeta_* = b(1 - \beta)C^{\beta-1}((C/C_*)^{mp-\beta} - 1). \quad (2.14)$$

Let  $p(m + \beta) \neq 1 + p$ . If  $C = C_*$ , then  $u_0$  is a stationary solution to (1.7), (1.10). If  $C \neq C_*$ , then the minimal solution to (1.7), (1.10) is of the self-similar form

$$u(x, t) = t^{1/(1-\beta)} f_1(\zeta), \zeta = xt^{-\frac{mp-\beta}{(1+p)(1-\beta)}}, \quad (2.15)$$

$$\eta(t) = \zeta_* t^{\frac{mp-\beta}{(1+p)(1-\beta)}}, 0 \leq t < +\infty, \quad (2.16)$$

where  $f_1(\zeta)$  solves the following nonlinear ODE problem

$$((f_1^m)'|^{p-1}(f_1^m)')' + \frac{mp-\beta}{(1+p)(1-\beta)} \zeta f_1' - \frac{1}{1-\beta} f_1 - b f_1^\beta = 0, \zeta \in \mathbb{R}, \quad (2.17)$$

$$f_1(\zeta) \sim C(-\zeta)^{(1+p)/(mp-\beta)}, \text{ as } \zeta \downarrow -\infty, \text{ and } f_1(\zeta) \rightarrow 0, \text{ as } \zeta \uparrow +\infty. \quad (2.18)$$

Moreover, there exists  $\zeta_*$  such that  $f(\zeta) \equiv 0$  for  $\zeta \geq \zeta_*$ ;  $f(\zeta) > 0$  for  $\zeta < \zeta_*$ . If  $C > C_*$

then the interface expands,  $f_1(0) = A_1 > 0$  (see Lemma 2.3.7), and

$$C_1 t^{\frac{1}{1-\beta}} (\zeta_1 - \zeta)_+^\mu \leq u \leq C_2 t^{\frac{1}{1-\beta}} (\zeta_2 - \zeta)_+^\mu, \quad 0 \leq x < +\infty, 0 < t < +\infty, \quad (2.19)$$

where

$$\begin{cases} \mu = p(mp-1)^{-1}, & \text{if } p(m+\beta) > 1+p, \\ \mu = (1+p)(mp-\beta)^{-1}, & \text{if } p(m+\beta) < 1+p, \end{cases}$$

which implies

$$\zeta_1 \leq \zeta_* \leq \zeta_2. \quad (2.20)$$

If  $0 < C < C_*$ , then the interface shrinks. If  $p(m+\beta) > 1+p$  then

$$\begin{aligned} & [C^{1-\beta}(-x)_+^{\frac{(1+p)(1-\beta)}{mp-\beta}} - b(1-\beta)t]_+^{\frac{1}{1-\beta}} \leq u \leq \\ & [C^{1-\beta}(-x)_+^{\frac{(1+p)(1-\beta)}{mp-\beta}} - b(1-\beta)(1-(C/C_*)^{mp-\beta})t]_+^{\frac{1}{1-\beta}}, \quad x \in \mathbb{R}, t \geq 0, \end{aligned} \quad (2.21)$$

which also implies (2.20), where we replace  $\zeta_1$  (respectively,  $\zeta_2$ ) with respective negative values given in Appendix 6.1. However, if  $p(m+\beta) < 1+p$ , then

$$C_* \left( -\zeta_3 t^{\frac{mp-\beta}{(1+p)(1-\beta)}} - x \right)_+^{\frac{1+p}{mp-\beta}} \leq u \leq C_3 \left( -\zeta_4 t^{\frac{mp-\beta}{(1+p)(1-\beta)}} - x \right)_+^{\frac{1+p}{mp-\beta}}, \quad x \in \mathbb{R}, t > 0, \quad (2.22)$$

where the left-hand side is valid for  $x \geq -\ell_0 t^{\frac{mp-\beta}{(1+p)(1-\beta)}}$ , while the right-hand side is valid for  $x \geq -\ell_1 t^{\frac{mp-\beta}{(1+p)(1-\beta)}}$ . From (2.22), (2.20) follows if we replace  $\zeta_1$  and  $\zeta_2$  with  $-\zeta_3$  and  $-\zeta_4$ , respectively.

*Further Details of Theorem 2.1.4.* There are four different subcases (see Figure 2.1).

(4a) If  $\beta = 1, \alpha = (1 + p)/(mp - 1)$ , the unique minimal solution to (1.7), (1.10) is

$$u_C = C(-x)_+^{(1+p)/(mp-1)} e^{-bt} [1 - (C/\bar{C})^{mp-1} b^{-1} (1 - e^{-b(mp-1)t})]^{1/(1-mp)} \quad (2.23)$$

where

$$T = +\infty, \quad \text{if } b \geq (C/\bar{C})^{mp-1},$$

$$T = (b(1 - mp))^{-1} \ln[1 - b(\bar{C}/C)^{mp-1}], \quad \text{if } -\infty < b < (C/\bar{C})^{mp-1},$$

(4b) Let  $\beta = 1$  and  $\alpha > (1 + p)/(mp - 1)$ . Then for any  $\varepsilon > 0$  there exists  $x_\varepsilon < 0$  and  $\delta_\varepsilon > 0$  such that

$$(C - \varepsilon)(-x)_+^\alpha e^{-bt} \leq u \leq (C + \varepsilon)(-x)_+^\alpha e^{-bt}$$

$$[1 - \varepsilon(b(mp - 1))^{-1} (1 - e^{-b(mp-1)t})]^{1/(1-mp)}, \quad x > x_\varepsilon, \quad 0 \leq t \leq \delta_\varepsilon. \quad (2.24)$$

(4c) Let  $1 < \beta < mp$  and  $\alpha \geq (1 + p)/(mp - \beta)$ . Then for any  $\varepsilon > 0$  there exists  $x_\varepsilon < 0$  and  $\delta_\varepsilon > 0$  such that

$$g_{-\varepsilon}(x, t) \leq u(x, t) \leq g_\varepsilon(x, t), \quad x \geq x_\varepsilon, \quad 0 \leq t \leq \delta_\varepsilon, \quad (2.25)$$

where

$$g_\varepsilon(x, t) = \begin{cases} [(C + \varepsilon)^{1-\beta} |x|^{\alpha(1-\beta)} + b(\beta - 1)(1 - \varepsilon - \kappa_\varepsilon)t]^{1/(1-\beta)}, & \text{if } x_\varepsilon \leq x < 0, \\ 0, & \text{if } x \geq 0, \end{cases}$$

where  $\kappa_\varepsilon = 0$ , if  $\alpha > (1 + p)/(mp - \beta)$ ;  $\kappa_\varepsilon = ((C + \varepsilon)/C_*)^{mp-\beta}$ , if  $\alpha = (1 + p)/(mp - \beta)$ .

(4d) Let either  $1 < \beta < mp$ ,  $(1 + p)/(mp - 1) \leq \alpha < (1 + p)/(mp - \beta)$ , or  $\beta \geq mp$ ,  $\alpha \geq$

$(1+p)/(mp-1)$ . If  $\alpha = (1+p)/(mp-1)$  then for any  $\varepsilon > 0$  there exists  $x_\varepsilon < 0$  and  $\delta_\varepsilon > 0$  such that

$$(C - \varepsilon)(-x)_+^{\frac{1+p}{mp-1}}(1 - \gamma_{-\varepsilon}t)^{\frac{1}{1-mp}} \leq u \leq (C + \varepsilon)(-x)_+^{\frac{1+p}{mp-1}}(1 - \gamma_\varepsilon t)^{\frac{1}{1-mp}} \quad (2.26)$$

for  $x > x_\varepsilon$ ,  $0 \leq t \leq \delta_\varepsilon$  (see 6.1 for  $\gamma_\varepsilon$ ). However, if  $\alpha > (1+p)/(mp-1)$ , then for arbitrary sufficiently small  $\varepsilon > 0$ , there exist  $x_\varepsilon < 0$  and  $\delta_\varepsilon > 0$  such that

$$(C - \varepsilon)(-x)_+^\alpha \leq u \leq (C + \varepsilon)(-x)_+^\alpha(1 - \varepsilon t)^{1/(1-mp)}, \quad x > x_\varepsilon, \quad 0 \leq t \leq \delta_\varepsilon. \quad (2.27)$$

*Results for the case  $b = 0$ .*

(1) If  $\alpha = p/(mp-1)$  the minimal solution to the problem (1.7), (1.10) is

$$u(x, t) = C(\xi_*t - x)_+^{p/(mp-1)}, \quad \xi_* = C^{mp-1} \left( \frac{mp}{mp-1} \right)^p. \quad (2.28)$$

If  $0 < \alpha < (1+p)/(mp-1)$ , then the minimal solution to (1.7), (1.10) has the self-similar form (2.8) and

$$\eta(t) = \xi_*t^{\frac{1}{1+p-\alpha(mp-1)}}, \quad 0 \leq t < +\infty, \quad (2.29)$$

where  $\xi_*$  and  $f$  solve (2.9)-(2.10). We have the estimation

$$C_4t^{\frac{\alpha}{1+p-\alpha(mp-1)}}(\xi_3 - \xi)_+^{\frac{p}{mp-1}} \leq u \leq C_5t^{\frac{\alpha}{1+p-\alpha(mp-1)}}(\xi_4 - \xi)_+^{\frac{p}{mp-1}}, \quad x \geq 0, \quad t \geq 0, \quad (2.30)$$

(see Appendix 6.1). If  $\alpha = p/(mp-1)$ , then  $\xi_3 = \xi_4 = \xi_*$  and both lower and upper estimations in (2.30) coincide with the solution (2.28).

(2) If  $\alpha = (1+p)/(mp-1)$ , then interface initially remains stationary. Explicit solu-

tion to (1.7), (1.10) is

$$u_C(x, t) = C(-x)_+^{(1+p)/(mp-1)} [\lambda(t_* - t)(1 - mp)]^{1/(1-mp)}, \quad x \in \mathbb{R}, \quad 0 \leq t < t_*, \quad (2.31)$$

where

$$t_* = 1/\lambda(1 - mp), \text{ with } \lambda = -C^{mp-1} \frac{p(m+1)(m(1+p))^p}{(mp-1)^{1+p}}.$$

(3) If  $\alpha > (1+p)/(mp-1)$ , then interfaces again remain stationary, and for any  $\varepsilon > 0$ , there exists a number  $x_\varepsilon < 0$  and  $\delta_\varepsilon > 0$ , such that

$$(C - \varepsilon)(-x)_+^\alpha \leq u \leq (C + \varepsilon)(-x)_+^\alpha (1 - \varepsilon t)^{1/(1-mp)}, \quad x_\varepsilon \leq x, \quad 0 \leq t \leq \delta_\varepsilon. \quad (2.32)$$

## 2.3 Preliminary Results

The following is a standard comparison result (comparison theorem) to be used throughout the dissertation.

**Lemma 2.3.1.** *Let  $g$  be a non-negative and continuous function in  $\overline{Q}$ , where:*

$$Q = \{(x, t) : \eta_0(t) < x < +\infty, \quad 0 < t < T \leq +\infty\}$$

$g = g(x, t)$  is in  $C_{x,t}^{2,1}$  in  $Q$  outside a finite number of curves:  $x = \eta_j(t)$ , which divide  $Q$  into a finite number of subdomains:  $Q^j$ , where  $\eta_j \in C[0, T]$ ; for arbitrary  $\delta > 0$  and finite  $\Delta \in (\delta, T]$  the function  $\eta_j$  is absolutely continuous in  $[\delta, \Delta]$ . Let  $g$  satisfy the inequality:

$$Lg \equiv g_t - \left( |(g^m)_x|^{p-1} (g^m)_x \right)_x + bg^\beta \geq 0, \quad (\leq 0),$$

at the points of  $Q$  where  $g \in C_{x,t}^{2,1}$ . Assume also that the function:  $|(g^m)_x|^{p-1} (g^m)_x$  is

continuous in  $Q$  and  $g \in L^\infty(Q \cap (t \leq T_1))$  for any finite  $T_1 \in (0, T)$ . If  $u$  is a weak solution of (1.7) in  $Q$ ,  $u \in C(\overline{Q})$ , and

$$g(\eta_0(t), t) \geq (\leq) u(\eta_0(t), t), \quad g(x, 0) \geq (\leq) u(x, 0).$$

Then

$$g \geq (\leq) u, \text{ in } \overline{Q}$$

Suppose that  $u_0 \in C(\mathbb{R})$ , and it has unbounded growth as  $|x| \rightarrow +\infty$ . It is well known that in this case some restriction must be imposed on the growth rate for existence, uniqueness of the solution to the CP (1.7), (1.8). For the particular cases of the equation (1.7) with  $b = 0$  this question was settled down in [44, 64] for the porous medium equation ( $p = 1$ ) with slow ( $m > 1$ ) and fast ( $0 < m < 1$ ) diffusion; and in [53, 54] for the  $p$ -Laplacian equation ( $m = 1$ ) with slow ( $p > 1$ ) and fast ( $0 < p < 1$ ) diffusion; The case of reaction-diffusion equation  $m > 1, p = 1, b > 0$  is analyzed in [71, 75, 26]. Surprisingly, only a partial result is available for the double-degenerate PDE (1.7). The sharp sufficient condition for the existence of the solution to the CP for (1.7),  $b = 0$  is established in [66]. In particular, it follows from [66] that the CP (1.7),(1.10) has a solution if and only if  $\alpha \leq (1 + p)/(mp - 1)$ . Moreover, solution is global ( $T = +\infty$ ) if  $\alpha < (1 + p)/(mp - 1)$  and only local in time if  $\alpha = (1 + p)/(mp - 1)$ . Uniqueness of the solution is an open problem. For our purposes it is satisfactory to employ the notion of the minimal solution.

**Definition 2.3.2** (Minimal Solution). Weak solution of the CP (1.7), (1.8) is called a *minimal solution* if

$$0 \leq u(x, t) \leq v(x, t), \tag{2.33}$$

for any weak solution  $v$  of the same problem (1.7), (1.8).

Note that the minimal solution is unique by definition. The following standard comparison result is true in the class of minimal solutions.

**Lemma 2.3.3.** *Let  $u$  and  $v$  be minimal solutions of the CP (1.7), (1.8). If*

$$u(x, 0) \geq (\leq) v(x, 0), \quad x \in \mathbb{R},$$

*then*

$$u(x, t) \geq (\leq) v(x, t), \quad (x, t) \in \mathbb{R} \times (0, T).$$

If the function  $u(x, t)$  is a minimal solution to CP (1.7), (1.10) with  $b = 0$ , then the function

$$\bar{u}(x, t) = \exp(-bt)u(x, (b(1 - mp))^{-1}(\exp(-b(mp - 1))t - 1)),$$

is a minimal solution to (1.7), (1.10) with  $b \neq 0$  and  $\beta = 1$ . Hence, from the above mentioned result it follows that the unique minimal solution to CP (1.7), (1.10) with  $mp > 1, b > 0, \beta = 1$ , and  $\alpha = (1 + p)/(mp - 1)$ , is the function  $\bar{u}_C(x, t)$  from (2.23).

### 2.3.1 Asymptotic Properties of Solutions With $mp > 1$

In the following lemmas we establish some preliminary estimations of the solution to the CP.

#### 2.3.1.1 Diffusion Dominates Reaction

**Lemma 2.3.4.** *If  $b = 0, 0 < \alpha < (1 + p)/(mp - 1)$ , then the minimal solution  $u$  of the CP (1.7), (1.10) has a self-similar form (2.8), where the self-similarity function  $f$  satisfies*

(2.11). If  $u_0$  satisfies (1.9), and  $u$  be a unique weak solution to CP (1.7), (1.8), then  $u$  satisfies (2.1)-(2.3).

*Proof of Lemma 2.3.4.* Let  $u$  be a unique minimal solution of the problem (1.7), (1.10).

If we consider a function

$$u_k(x, t) = ku(k^{-1/\alpha}x, k^{(\alpha(mp-1)-(1+p))/\alpha}t), \quad k > 0, \quad (2.34)$$

it may easily be checked that this satisfies (1.7), (1.10). Since  $u$  is a minimal solution we have

$$u(x, t) \leq ku(k^{-1/\alpha}x, k^{(\alpha(mp-1)-(1+p))/\alpha}t), \quad k > 0. \quad (2.35)$$

By changing the variable in (4.38) as

$$y = k^{-1/\alpha}x, \quad \tau = k^{(\alpha(mp-1)-(1+p))/\alpha}t, \quad (2.36)$$

we derive (4.38) with  $k$  replaced with  $k^{-1}$ . Since  $k > 0$  is arbitrary, (4.38) follows with "=". If we choose  $k = t^{\alpha/(1+p-\alpha(mp-1))}$ , the latter implies (2.8) with  $f(\xi) = u(\xi, 1)$ , where  $f$  is a nonnegative and continuous solution of (2.9), (2.10). By [42], PDE (1.7) has a finite speed of propagation property, and minimal solution of (1.7), (1.10) has an expanding interface. Therefore, the upper bound  $\xi_*$  of the support of  $f$  is positive and finite;  $f$  is positive and smooth for  $\xi < \xi_*$  and  $f = 0$  for  $\xi \geq \xi_*$ . Thus, (2.29) is valid. Proof of (2.2) and (2.11) coincide with the proof given in Lemma 3.2 of [23].

Now suppose that  $u_0$  satisfies (1.9). Then for arbitrary sufficiently small  $\varepsilon > 0$ , there exists an  $x_\varepsilon < 0$  such that

$$(C - \varepsilon)(-x)_+^\alpha \leq u_0(x) \leq (C + \varepsilon)(-x)_+^\alpha, \quad x \geq x_\varepsilon. \quad (2.37)$$



Let  $u_\varepsilon(x, t)$  (respectively,  $u_{-\varepsilon}(x, t)$ ) be a minimal solution to the CP (1.7), (1.8) with initial data  $(C + \varepsilon)(-x)_+^\alpha$  (respectively,  $(C - \varepsilon)(-x)_+^\alpha$ ). Since the solution to the CP (1.7), (1.8) is continuous, there exists a number  $\delta = \delta(\varepsilon) > 0$  such that

$$u_\varepsilon(x_\varepsilon, t) \geq u(x_\varepsilon, t), \quad u_{-\varepsilon}(x_\varepsilon, t) \leq u(x_\varepsilon, t), \quad \text{for } 0 \leq t \leq \delta. \quad (2.38)$$

From (2.37), (2.38), and by applying the comparison result, lemma 2.3.1, it follows that:

$$u_{-\varepsilon} \leq u \leq u_\varepsilon, \quad \text{for } x \geq x_\varepsilon, \quad 0 \leq t \leq \delta. \quad (2.39)$$

We have

$$u_{\pm\varepsilon}(\xi_\rho(t), t) = t^{\alpha/(1+p-\alpha(mp-1))} f(\rho; C \pm \varepsilon), \quad \rho < \xi_*, \quad t \geq 0. \quad (2.40)$$

(Furthermore, we denote the right-hand side of (2.11a) by  $f(\rho, C)$ ). Now taking  $x = \xi_\rho(t)$  in (2.39), after multiplying by  $t^{-\alpha/(1+p-\alpha(mp-1))}$  and passing to the limit, first as  $t \rightarrow 0^+$  and then as  $\varepsilon \rightarrow 0^+$ , we can easily derive (2.3). Similarly, from (2.39), (2.40) and (2.29), (2.1) easily follows. The lemma is proved.  $\square$

**Lemma 2.3.5.** *If either  $b > 0$  and  $0 < \alpha < \frac{1+p}{mp-\min\{1,\beta\}}$  or  $b < 0$ ,  $\beta \geq 1$  and  $0 < \alpha < \frac{1+p}{mp-1}$  and  $u$  is the unique weak solution to the CP (1.7)-(1.9), then  $u$  satisfies (2.3).*

*Proof of Lemma 2.3.5.* Let  $b > 0$ . Then, as in the proof of Lemma 2.3.4, (2.37) and (2.38) follow from (1.9). From [66] and Lemma 2.3.3 it follows that the existence, uniqueness, and comparison result for the minimal solution of the CP (1.7), (1.8) with  $u_0 = (C \pm \varepsilon)(-x)_+^\alpha$  and  $T = +\infty$  hold. As before, from (2.37) and (2.38), (2.39) follows. For arbitrary  $k > 0$ , the function

$$u_k^{\pm\varepsilon}(x, t) = k u_{\pm\varepsilon} \left( k^{-1/\alpha} x, k^{(\alpha(mp-1)-(1+p))/\alpha} t \right), \quad k > 0, \quad (2.41)$$

is a minimal solution of the following problem

$$u_t - \left( |(u^m)_x|^{p-1} (u^m)_x \right)_x + bk^{(\alpha(mp-\beta)-(1+p))/\alpha} u^\beta = 0, \quad x \in \mathbb{R}, t > 0, \quad (2.42a)$$

$$u(x, 0) = (C \pm \varepsilon)(-x)_+^\alpha, \quad x \in \mathbb{R}. \quad (2.42b)$$

Since  $\alpha(mp - \beta) - (1 + p) < 0$ , it follows that

$$\lim_{k \rightarrow +\infty} u_k^{\pm \varepsilon}(x, t) = v_{\pm \varepsilon}(x, t), \quad x \in \mathbb{R}, t \geq 0, \quad (2.43)$$

where  $v_{\pm \varepsilon}$  is a minimal solution to CP (1.7), (1.8) with  $b = 0$ ,  $u_0 = (C \pm \varepsilon)(-x)_+^\alpha$ , and  $T = +\infty$ . Hence,  $v_{\pm \varepsilon}$  satisfies (2.40). Taking  $x = \xi_\rho(t)$ , where  $\rho < \xi_*$  is fixed, from (2.43) it follows that for arbitrary  $t > 0$

$$\lim_{k \rightarrow +\infty} k u_{\pm \varepsilon} \left( k^{-1/\alpha} \xi_\rho(t), k^{(\alpha(mp-1)-(1+p))/\alpha} t \right) = t^{\alpha(1+p-\alpha(mp-1))} f(\rho; C \pm \varepsilon). \quad (2.44)$$

Letting  $\tau = k^{(\alpha(mp-1)-(1+p))/\alpha} t$ , then (2.44) implies

$$u_{\pm \varepsilon}(\xi_\rho(\tau), \tau) \sim \tau^{\alpha/(1+p-\alpha(mp-1))} f(\rho; C \pm \varepsilon), \quad \text{as } \tau \rightarrow 0^+. \quad (2.45)$$

As before, (2.3) easily follows from (2.39) and (2.45).

Let  $b < 0$ . From (1.9) it follows that for all  $\varepsilon > 0$  there exists a number  $x_\varepsilon < 0$  such that

$$(C - \varepsilon/2)(-x)_+^\alpha \leq u_0(x) \leq (C + \varepsilon/2)(-x)_+^\alpha, \quad x \geq x_\varepsilon. \quad (2.46)$$

Assume that  $u_{\pm \varepsilon}$  is a solution of the boundary value problem

$$Lu_{\pm \varepsilon} = 0, \quad |x| < |x_\varepsilon|, \quad 0 < t \leq \delta, \quad (2.47)$$

$$u_{\pm\varepsilon}(x, 0) = (C \pm \varepsilon)(-x)_+^\alpha, \quad |x| \leq |x_\varepsilon|,$$

$$u_{\pm\varepsilon}(x_\varepsilon, t) = (C \pm \varepsilon)(-x_\varepsilon)^\alpha, \quad u_{\pm\varepsilon}(-x_\varepsilon, t) = u(-x_\varepsilon, t) = 0, \quad 0 \leq t \leq \delta, \quad (2.48)$$

where  $\delta = \delta(\varepsilon) > 0$  is chosen such that

$$u_\varepsilon(x_\varepsilon, t) \geq u(x_\varepsilon, t), \quad u_{-\varepsilon}(x_\varepsilon, t) \leq u(x_\varepsilon, t), \quad 0 \leq t \leq \delta. \quad (2.49)$$

From the comparison theorem it follows that

$$u_{-\varepsilon} \leq u \leq u_\varepsilon, \quad |x| \leq |x_\varepsilon|, \quad 0 \leq t \leq \delta. \quad (2.50)$$

Now if we rescale

$$u_k^{\pm\varepsilon}(x, t) = k u_{\pm\varepsilon}(k^{-1/\alpha} x, k^{(\alpha(mp-1)-(1+p))/\alpha} t), \quad k > 0, \quad (2.51)$$

then  $u_k^{\pm\varepsilon}$  satisfies the following problem

$$L_k u_k^{\pm\varepsilon} \equiv (u_k^{\pm\varepsilon})_t - \left( |(u_k^{\pm\varepsilon})_x|^{p-1} ((u_k^{\pm\varepsilon})_x)_x \right) + b k^{(\alpha(mp-\beta)-(1+p))/\alpha} (u_k^{\pm\varepsilon})^\beta = 0 \quad \text{in } D_\varepsilon^k, \quad (2.52)$$

$$u_k^{\pm\varepsilon}(x, 0) = (C \pm \varepsilon)(-x)_+^\alpha, \quad |x| \leq k^{1/\alpha} |x_\varepsilon|, \quad (2.53)$$

$$u_k^{\pm\varepsilon}(k^{1/\alpha} x_\varepsilon, t) = k(C \pm \varepsilon)(-x_\varepsilon)^\alpha, \quad u_k^{\pm\varepsilon}(-k^{1/\alpha} x_\varepsilon, t) = 0, \quad 0 \leq t \leq k^{\frac{p+1+\alpha(1-mp)}{\alpha}} \delta, \quad (2.54)$$

where

$$D_\varepsilon^k = \{(x, t) : |x| < k^{1/\alpha}|x_\varepsilon|, 0 < t \leq k^{(\alpha(1-mp)+(1+p))/\alpha}\delta\}.$$

The next step is to prove the convergence of the sequence  $\{u_k^{\pm\varepsilon}\}$  as  $k \rightarrow +\infty$ . If  $b > 0$  the proof coincides with that given for Lemma 3.2 of [23]. If  $b < 0$  we consider a function

$$g(x, t) = (C + 1)(1 + |x|^\mu)^{\frac{\alpha}{\mu}}(1 - vt)^\gamma, \quad x \in \mathbb{R}, 0 \leq t \leq t_0 = v^{-1}/2,$$

where

$$\begin{aligned} \gamma < 0, \mu > \frac{p+1}{p}, v = -h_* + 1, h_* = \min_{\mathbb{R}} h(x) > -\infty \\ h(x) = p(\alpha m)^p (C + 1)^{p-1} \gamma^{-1} (1 - vt)^{\gamma(mp-1)+1} (1 + |x|^\mu)^{\frac{\alpha(mp-1)-\mu(p+1)}{\mu}} |x|^{(\mu-1)p-1} \\ \times [(\mu-1)(1 + |x|^\mu) + (\alpha m - \mu)\mu|x|^\mu]. \end{aligned}$$

Then we have

$$\begin{aligned} L_k g &= -\gamma(C + 1)(1 + |x|^\mu)^{\frac{\alpha}{\mu}}(1 - vt)^{\gamma-1} S \quad \text{in } D_\varepsilon^k, \\ S &= v + h(x) - b(C + 1)^{\beta-1} \gamma^{-1} (1 + |x|^\mu)^{\frac{\alpha(\beta-1)}{\mu}} (1 - vt)^{\gamma(\beta-1)+1} k^{\frac{\alpha(mp-\beta)-(1+p)}{\alpha}}, \end{aligned}$$

and therefore

$$S \geq 1 + B, \quad \text{in } D_{0\varepsilon}^k = D_\varepsilon^k \cap \{(x, t) : 0 < t \leq t_0\}, \quad (2.55)$$

where

$$B = O(k^{mp-1-(1+p)/\alpha}) \quad \text{uniformly for } (x, t) \in D_{0\varepsilon}^k \text{ as } k \rightarrow +\infty.$$

Hence, we have for  $0 < \varepsilon \ll 1$  and  $k \gg 1$

$$L_k g \geq 0, \text{ in } D_{0\varepsilon}^k, \quad (2.56)$$

$$g(x, 0) \geq u_k^{\pm\varepsilon}(x, 0), \text{ for } |x| \leq k^{1/\alpha}|x_\varepsilon|, \quad (2.57)$$

$$g(\pm k^{1/\alpha}x_\varepsilon, t) \geq u_k^{\pm\varepsilon}(\pm k^{1/\alpha}x_\varepsilon, t), \text{ for } 0 \leq t \leq t_0. \quad (2.58)$$

From (2.56)-(2.58) and comparison theorem it follows that

$$0 \leq u_k^{\pm\varepsilon}(x, t) \leq g(x, t), \text{ in } \bar{D}_{0\varepsilon}^k \quad (2.59)$$

Let  $G$  be an arbitrary fixed compact subset of

$$P = \{(x, t) : x \in \mathbb{R}, 0 < t \leq t_0\}$$

By choosing  $k$  to be so large that  $G \subset P$ , it follows from (2.59) that the sequences  $\{u_k^{\pm\varepsilon}\}$  are uniformly bounded in  $G$ . From [68, 67], it follows that they are uniformly Hölder continuous in  $G$ . From the Arzelá-Ascoli theorem and standard diagonalization argument it follows that there exist functions  $v_{\pm\varepsilon}$  such that for some subsequence  $k'$

$$\lim_{k' \rightarrow +\infty} u_{k'}^{\pm\varepsilon}(x, t) = v_{\pm\varepsilon}(x, t), \quad (x, t) \in P. \quad (2.60)$$

It may be easily checked that  $v_{\pm\varepsilon}$  is a solution of the CP (1.7), (1.8) with  $u_0 = (C \pm \varepsilon)(-x)_+^\alpha$ .

The lemma is proved. □

### 2.3.1.2 Diffusion and Reaction are in Balance

In the next lemma we analyze a special class of finite travelling wave solutions. By a finite travelling-wave solution with velocity  $0 \neq k \in \mathbb{R}$  we mean a solution  $u(x, t) = \varphi(kt - x)$ , where  $\varphi(y) \geq 0$ ,  $\varphi \not\equiv 0$ , and  $\varphi(y) = 0$  for  $y \leq y_0$  for some  $y_0 \in \mathbb{R}$ .

**Lemma 2.3.6.** *Let  $0 < \beta < 1$ ,  $p(m + \beta) > 1 + p$ . PDE (1.7) admits a finite travelling-wave solution,  $u(x, t) = \varphi(kt - x)$ , with  $\varphi(y) = 0$  for  $y \leq 0$ ;  $\varphi(y) > 0$ , for  $y > 0$ , and*

$$\lim_{y \rightarrow +\infty} y^{-\frac{1+p}{mp-\beta}} \varphi(y) = C_*. \quad (2.61)$$

The formula (2.61) is formula 2. from Theorem 3.1.1 of Chapter 3. Please refer to the proof of Theorem 3.1.1 for the proof of (2.61).

**Lemma 2.3.7.** *If  $0 < \beta < 1$ ,  $\alpha = (1 + p)/(mp - \beta)$ , then the minimal solution  $u$  to the CP (1.7), (1.10) has a self-similar form (2.15), where the self-similarity function  $f$  satisfies (4.34), (2.18). If  $C > C_*$ , then  $f_1(0) = A_1 > 0$ , where  $A_1$  depends on  $m, p, \beta, C$  and  $b$ . If  $u_0$  satisfies (1.9) with  $\alpha = (1 + p)/(mp - \beta)$ ,  $C > C_*$ , and  $u$  be a unique weak solution to CP (1.7), (1.8), then  $u$  satisfies*

$$u(0, t) \sim A_1 t^{1/(1-\beta)}, \text{ as } t \rightarrow 0^+ \quad (2.62)$$

*Proof of Lemma 2.3.7.* Let  $u$  be a unique minimal solution of the problem (1.7), (1.10) [66, 92]. Rescaled function

$$u_k(x, t) = ku(k^{-1/\alpha}x, k^{\beta-1}t), \quad k > 0, \quad (2.63)$$

satisfies (1.7), (1.10), and therefore

$$u(x, t) \leq ku(k^{-1/\alpha}x, k^{\beta-1}t), \quad k > 0. \quad (2.64)$$

As in the proof of lemma 2.3.4, it follows that (2.64) is true with equality sign. If we choose  $k = t^{1/(1-\beta)}$ , then (2.64) implies (2.15) with  $f_1(\zeta) = u(\zeta, 1)$ , where  $f_1$  is a nonnegative and continuous solution of (4.34), (2.18). By [42], PDE (1.7) has a finite speed of propagation property, and minimal solution of (1.7), (1.10) has a finite interface. Therefore, upper bound  $\xi_*$  of the support of  $f$  is finite;  $f$  is positive and smooth for  $\xi < \xi_*$  and  $f = 0$  for  $\xi \geq \xi_*$ . Now we prove that if  $C > C_*$ , then  $f_1(0) = A_1 > 0$ . We divide the proof into two cases:

**Case 1:**  $p(m + \beta) < 1 + p$

It is enough to show that there exists  $t_0 > 0$  such that  $u(0, t_0) > 0$ . Let  $g(x, t) = C_1(t - x)_+^{\frac{1+p}{mp-\beta}}$ ,  $C_1 \in (C_*, C)$ .

$$Lg = bC_1^\beta(t-x)_+^{\frac{\beta(1+p)}{mp-\beta}} \left[ 1 - \left( \frac{C_1}{C_*} \right)^{mp-\beta} + C_1^{1-\beta} \frac{1+p}{b(mp-\beta)} (t-x)_+^{\frac{1+p-p(m+\beta)}{mp-\beta}} \right]$$

Since  $C_1 < C$ , we can choose  $x_1 < 0$  and  $\delta > 0$  such that

$$Lg \leq 0, \text{ in } Q := \{(x, t) : x_1 \leq x < t, 0 < t \leq \delta\},$$

$$g(x, 0) \leq u(x, 0), \quad x_1 \leq x; \quad g(x_1, t) \leq u(x_1, t), \quad 0 \leq t \leq \delta.$$

Comparison Lemma 2.3.1 implies

$$0 < g(x, t) \leq u(x, t), \quad x_1 \leq x < t, \quad 0 \leq t \leq \delta.$$

In particular, we have:  $u(0, t_0) > 0$  for all  $0 < t_0 \leq \delta$ , which implies that  $f_1(0) = A_1 > 0$ .

**Case 2:**  $p(m + \beta) > 1 + p$

We apply Lemma 2.3.6 with the forward traveling wave ( $k > 0$ ). By (2.61) for some  $M > 0$  we have

$$\varphi(y) < Cy^{\frac{1+p}{mp-\beta}}, \text{ for } y > M. \quad (2.65)$$

We choose

$$K = \max\{\varphi(y) : 0 \leq y \leq M\}, \quad \xi = \max\left\{M; \left(\frac{K}{C}\right)^{\frac{mp-\beta}{1+p}}\right\}, \quad (2.66)$$

and consider a family of traveling-wave solutions to (1.7) of the form:  $g(x, t) = \varphi(kt - x - \xi)$ . From (2.65), (2.66) it follows that

$$\varphi(-x - \xi) \leq C(-x)_+^{\frac{1+p}{mp-\beta}}, \text{ for any } x \in \mathbb{R}. \quad (2.67)$$

From the comparison theorem it follows that  $g \leq u$  for any  $x \in \mathbb{R}, t \geq 0$ . By choosing  $t_0 > 0$  such that  $k > \xi t_0^{-1} > 0$ , we ensure that

$$0 < g(0, t_0) = \varphi(kt_0 - \xi) \leq u(0, t_0) = t_0^{\frac{1}{1-\beta}} f_1(0), \quad (2.68)$$

which proves that  $f_1(0) > 0$ . To prove the asymptotic formula (2.62) we proceed as we did in the proof in Lemma 2.3.5. As before, (2.37)-(2.39) follow from (1.9), where  $v_{\pm\epsilon}$  is a solution of the problem

$$v_t - \left(|(v^m)_x|^{p-1}(v^m)_x\right)_x + bv^\beta = 0, \quad |x| < |x_\epsilon|, \quad 0 < t \leq \delta, \quad (2.69)$$

$$v(x_\epsilon, t) = (C \pm \epsilon)(-x_\epsilon)^\alpha, \quad v(-x_\epsilon, t) = u(-x_\epsilon, t), \quad 0 \leq t \leq \delta, \quad (2.70)$$

$$v(x, 0) = (C \pm \epsilon)(-x)_+^\alpha, \quad |x| \leq |x_\epsilon|. \quad (2.71)$$



Rescaled function

$$u_k^{\pm\varepsilon}(x, t) = ku_{\pm\varepsilon}\left(k^{-\frac{1}{\alpha}}x, k^{\beta-1}t\right), \quad k > 0,$$

satisfies the Dirichlet problem:

$$v_t = \left(|(v^m)_x|^{p-1}(v^m)_x\right)_x - bv^\beta, \quad \text{in } E_\varepsilon^k = \{|x| < k^{\frac{1}{\alpha}}|x_\varepsilon|, 0 < t \leq k^{1-\beta}\delta\}, \quad (2.72a)$$

$$v(k^{\frac{1}{\alpha}}x_\varepsilon, t) = k(C \pm \varepsilon)(-x_\varepsilon)^\alpha, \quad v(-k^{\frac{1}{\alpha}}x_\varepsilon, t) = ku(-x_\varepsilon, k^{\beta-1}t), \quad 0 \leq t \leq k^{1-\beta}\delta, \quad (2.72b)$$

$$v(x, 0) = (C \pm \varepsilon)(-x)_+^\alpha, \quad |x| \leq k^{\frac{1}{\alpha}}|x_\varepsilon|. \quad (2.72c)$$

As before in the proof of Lemma 2.3.5 we have:

$$\lim_{k \rightarrow +\infty} u_k^{\pm\varepsilon}(x, t) = v_{\pm\varepsilon}(x, t), \quad (x, t) \in P := \{(x, t) : x \in \mathbb{R}, 0 < t \leq t_0\}, \quad (2.73)$$

thus,

$$v_{\pm\varepsilon}(x, t) = t^{\frac{1}{1-\beta}} f_1(\rho; C \pm \varepsilon), \quad \rho < \zeta_*, \quad t \geq 0 \quad (2.74)$$

Taking  $x = \eta_\rho(t) = \rho t^{\frac{mp-\beta}{(1+p)(1-\beta)}}$  and  $\tau = k^{\beta-1}t$  it follows from (2.73) that

$$u_{\pm\varepsilon}(\eta_\rho(\tau), \tau) \sim \tau^{\frac{1}{1-\beta}} f_1(\rho; C \pm \varepsilon), \quad \text{as } \tau \rightarrow 0^+. \quad (2.75)$$

From (2.39) and (2.75), since  $\varepsilon > 0$  is arbitrary and  $f_1(0) = A_1 > 0$ , the desired asymptotic formula (2.62) follows. The lemma is proved.  $\square$

### 2.3.1.3 Reaction Dominates Diffusion

**Lemma 2.3.8.** *If  $0 < \beta < 1$ ,  $\alpha > (1+p)/(mp-\beta)$ , and  $u$  be a unique weak solution to CP (1.7)-(1.9), then  $u$  satisfies (2.7).*

*Proof of Lemma 2.3.8.* As before, (2.37)-(2.39) follow from (1.9), where  $v_{\pm\varepsilon}$  is a solu-

tion of the problem

$$v_t - \left( |(v^m)_x|^{p-1} (v^m)_x \right)_x + bv^\beta = 0, \quad |x| < |x_\varepsilon|, \quad 0 < t \leq \delta, \quad (2.76)$$

$$v(x_\varepsilon, t) = (C \pm \varepsilon)(-x_\varepsilon)^\alpha, \quad v(-x_\varepsilon, t) = u(-x_\varepsilon, t), \quad 0 \leq t \leq \delta, \quad (2.77)$$

$$v(x, 0) = (C \pm \varepsilon)(-x)_+^\alpha, \quad |x| \leq |x_\varepsilon|. \quad (2.78)$$

Rescaled function

$$u_k^{\pm\varepsilon}(x, t) = ku_{\pm\varepsilon}\left(k^{-\frac{1}{\alpha}}x, k^{\beta-1}t\right), \quad k > 0,$$

satisfies the Dirichlet problem

$$v_t = k^{\frac{1+p-\alpha(mp-\beta)}{\alpha}} \left( |(v^m)_x|^{p-1} (v^m)_x \right)_x - bu^\beta, \quad \text{in } E_\varepsilon^k, \quad (2.79a)$$

$$v(k^{\frac{1}{\alpha}}x_\varepsilon, t) = k(C \pm \varepsilon)(-x_\varepsilon)^\alpha, \quad v(-k^{\frac{1}{\alpha}}x_\varepsilon, t) = ku(-x_\varepsilon, k^{\beta-1}t), \quad 0 \leq t \leq k^{1-\beta}\delta, \quad (2.79b)$$

$$v(x, 0) = (C \pm \varepsilon)(-x)_+^\alpha, \quad |x| \leq k^{\frac{1}{\alpha}}|x_\varepsilon|, \quad (2.79c)$$

where

$$E_\varepsilon^k := \left\{ |x| < k^{\frac{1}{\alpha}}|x_\varepsilon|, \quad 0 < t \leq k^{1-\beta}\delta \right\}.$$

The next step consists in proving the convergence of the sequence  $\{u_k^{\pm\varepsilon}\}$  as  $k \rightarrow +\infty$ . This step is identical with the proof given in the similar Lemma 3.4 from [23]. For any fixed  $t_0 > 0$ , the function  $g(x, t) = (C + 1)(1 + x^2)^{\alpha/2}e^t$  is a uniform upper bound for the sequence  $\{u_k^{\pm\varepsilon}\}$  in  $E_{0\varepsilon}^k = E_\varepsilon^k \cap P$ , where  $P = \{(x, t) : 0 < t \leq t_0\}$ . The sequences  $\{u_k^{\pm\varepsilon}\}$  are uniformly Hölder continuous on an arbitrary compact subset of  $P$  [57, 68]. As in the proof of the Lemma 3.4 of [23] it is proved that some subsequences  $\{u_{k'}^{\pm\varepsilon}\}$  converge to

solutions of the reaction equation. This implies

$$u_{\pm\varepsilon}(\eta_\ell(\tau), \tau) \sim \tau^{\frac{1}{1-\beta}} \left[ (C \pm \varepsilon)^{1-\beta} \ell^{\alpha(1-\beta)} - b(1-\beta) \right]^{\frac{1}{1-\beta}}, \text{ as } \tau \rightarrow 0^+. \quad (2.80)$$

From (2.39) and (2.80), since  $\varepsilon > 0$  is arbitrary, the desired formula (2.7) follows. The lemma is proved.  $\square$

## 2.4 Proofs of the Main Results

### 2.4.1 Proof of Theorem 2.1.1

*Proof of Theorem 2.1.1.* From Lemma 2.3.5 and (2.3) it follows

$$\liminf_{t \rightarrow 0^+} \eta(t) t^{1/(\alpha(mp-1)-(1+p))} \geq \xi_*. \quad (2.81)$$

For any  $\varepsilon > 0$ , let  $u_\varepsilon$  be a minimal solution of the CP (1.7), (1.10) with  $b = 0$  and with  $C$  replaced by  $C + \varepsilon$ . The second inequality of (2.37) and the first inequality of (2.38) follow from (1.9). Since  $u_\varepsilon$  is a supersolution of (1.7), from (2.37), (2.38), and a comparison principle, the second inequality of (2.39) follows. By Lemma 2.3.4 we have

$$\eta(t) \leq (C + \varepsilon)^{\frac{mp-1}{1+p-\alpha(mp-1)}} \xi_*' t^{1/(1+p-\alpha(mp-1))}, \quad 0 \leq t \leq \delta,$$

and hence

$$\limsup_{t \rightarrow 0^+} \eta(t) t^{1/(\alpha(mp-1)-(1+p))} \leq \xi_*. \quad (2.82)$$

From (2.81) and (2.82), (2.1) follows.  $\square$

## 2.4.2 Proof of Theorem 2.1.2

*Proof of Theorem 2.1.2.* Assume that  $u_0$  is defined by (1.10) and  $p(m+\beta) \neq 1+p$ . The self-similar form (2.15) follows from Lemma 2.3.7. Let  $C > C_*$ . For a function

$$g(x, t) = t^{1/(1-\beta)} f_1(\zeta), \quad \zeta = xt^{-\frac{mp-\beta}{(1+p)(1-\beta)}}. \quad (2.83)$$

we have

$$Lg = t^{\frac{\beta}{1-\beta}} \mathcal{L}^0 f_1, \quad (2.84)$$

where the operator  $\mathcal{L}^0$  is defined by (4.34). By choosing

$$f_1(\zeta) = C_0(\zeta_0 - \zeta)_+^{\gamma_0}, \quad 0 < \zeta < +\infty,$$

with  $C_0, \zeta_0 > 0$  and  $\gamma_0 = (1+p)/(mp-\beta)$  we have

$$\mathcal{L}^0 f_1 = bC_0^\beta (\zeta_0 - \zeta)_+^{\frac{\beta(1+p)}{mp-\beta}} \left\{ 1 - (C_0/C_*)^{mp-\beta} + \frac{C_0^{1-\beta}}{b(1-\beta)} \zeta_0 (\zeta_0 - \zeta)_+^{\frac{1+p-p(m+\beta)}{mp-\beta}} \right\}. \quad (2.85)$$

For an upper estimation we choose  $C_0 = C_2$  and  $\zeta_0 = \zeta_2$  (see Appendix 6.1). If  $p(m+\beta) > 1+p$ , we have

$$\mathcal{L}^0 f_1 \geq bC_2^\beta (\zeta_2 - \zeta)_+^{\frac{\beta(1+p)}{mp-\beta}} \left\{ 1 - (C_2/C_*)^{mp-\beta} + \frac{C_2^{1-\beta}}{b(1-\beta)} \zeta_2^{\frac{(1+p)(1-\beta)}{mp-\beta}} \right\} = 0, \quad \text{for } 0 \leq \zeta \leq \zeta_2,$$

while if  $p(m+\beta) < 1+p$ , we have

$$\mathcal{L}^0 f_1 \geq bC_2^\beta (\zeta_2 - \zeta)_+^{\frac{\beta(1+p)}{mp-\beta}} \left\{ 1 - (C_2/C_*)^{mp-\beta} \right\} = 0, \quad \text{for } 0 \leq \zeta \leq \zeta_2.$$

By (4.44) we have

$$Lg \geq 0, \text{ for } 0 < x < \zeta_2 t^{\frac{mp-\beta}{(1+p)(1-\beta)}}, 0 < t < +\infty, \quad (2.86a)$$

$$Lg = 0, \text{ for } x > \zeta_2 t^{\frac{mp-\beta}{(1+p)(1-\beta)}}, 0 < t < +\infty. \quad (2.86b)$$

Lemma 2.3.1 implies that  $g$  is a supersolution of (1.7) in  $\{(x, t) : x > 0, t > 0\}$ . Since

$$g(x, 0) = u(x, 0) = 0, \text{ for } 0 \leq x < +\infty, \quad (2.87a)$$

$$g(0, t) = u(0, t), \text{ for } 0 \leq x < +\infty, \quad (2.87b)$$

the right-hand side of (2.19) follows. If  $p(m+\beta) < 1+p$ , to prove the lower estimation we choose  $C_0 = C_1$ ,  $\zeta_0 = \zeta_1$ , and  $\gamma_0 = (1+p)/(mp-\beta)$ . From (2.85) and (4.44) we have

$$\mathcal{L}^0 f_1 \leq bC_1^\beta (\zeta_1 - \zeta)^{\frac{\beta(1+p)}{mp-\beta}} \left\{ 1 - (C_1/C_*)^{mp-\beta} + \frac{C_1^{1-\beta}}{b(1-\beta)} \zeta_1^{\frac{(1+p)(1-\beta)}{mp-\beta}} \right\} = 0, \text{ for } 0 \leq \zeta \leq \zeta_1,$$

$$Lg \leq 0, \text{ for } 0 < x < \zeta_1 t^{\frac{mp-\beta}{(1+p)(1-\beta)}}, 0 < t < +\infty, \quad (2.88a)$$

$$Lg = 0, \text{ for } x > \zeta_1 t^{\frac{mp-\beta}{(1+p)(1-\beta)}}, 0 < t < +\infty. \quad (2.88b)$$

As before from (2.87) and Lemma 2.3.1, the left-hand side of (2.19) follows. If  $p(m+\beta) > 1+p$ , then to prove the lower estimation we choose  $C_0 = C_1$ ,  $\zeta_0 = \zeta_1$  and  $\gamma_0 =$

$p/(mp-1)$ . We have

$$\begin{aligned} \mathcal{L}^0 f_1 &\leq C_1(1-\beta)^{-1}(\zeta_1 - \zeta)^{\frac{1+p-mp}{mp-1}} \times \\ &\times \left\{ \zeta_1 - C_1^{mp-1} \frac{(1-\beta)p(mp)^p}{(mp-1)^{1+p}} + b(1-\beta)C_1^{\beta-1} \zeta_1^{\frac{p(m+\beta)-(1+p)}{mp-1}} \right\} = 0, \quad 0 < \zeta < \zeta_1, \end{aligned}$$

which again implies (2.88). From Lemma 2.3.1, the left-hand side of (2.19) follows.

Let  $p(m+\beta) > 1+p$  and  $0 < C < C_*$ . For  $\gamma \in [0, 1)$  consider a function

$$g(x, t) = \left[ C^{1-\beta}(-x)_+^{\frac{(1+p)(1-\beta)}{mp-\beta}} - b(1-\beta)(1-\gamma)t \right]_+^{\frac{1}{1-\beta}}, \quad x \in \mathbb{R}, \quad t > 0.$$

We estimate  $Lg$  in

$$M := \{(x, t) : -\infty < x < \mu_\gamma(t), \quad t > 0\}, \quad \mu_\gamma(t) = -\left[ b(1-\beta)(1-\gamma)C^{\beta-1}t \right]^{\frac{mp-\beta}{(1+p)(1-\beta)}}.$$

We have  $Lg = bg^\beta S$ , where

$$\begin{aligned} S &= \gamma - C^{mp-\beta} \left[ 1 - \left( \frac{-\mu_\gamma(t)}{(-x)_+} \right)^{\frac{(1+p)(1-\beta)}{mp-\beta}} \right]^{\frac{p(m+\beta-1)-\beta}{1-\beta}} \times \\ &\times \left[ R_1 + R_2 \left[ 1 - \left( \frac{-\mu_\gamma(t)}{(-x)_+} \right)^{\frac{(1+p)(1-\beta)}{mp-\beta}} \right]^{-1} \right], \end{aligned} \quad (2.89a)$$

$$S|_{t=0} = \gamma - (C/C_*)^{mp-\beta}, \quad S|_{x=\mu_\gamma(t)} = \gamma, \quad (2.89b)$$

where  $R_1, R_2 > 0$  (see Appendix 6.1). Moreover,

$$S_t \geq 0 \text{ in } M.$$

Thus,

$$\gamma - (C/C_*)^{mp-\beta} \leq S \leq \gamma \text{ in } M.$$

If we take  $\gamma = (C/C_*)^{mp-\beta}$  (respectively,  $\gamma = 0$ ), then we have

$$Lg \geq 0 \text{ (respectively, } Lg \leq 0) \text{ in } M, \quad (2.90a)$$

$$Lg = 0, \text{ for } x > \mu_\gamma(t), t > 0. \quad (2.90b)$$

From Lemma 2.3.1, the estimation (2.21) follows. Let  $p(m+\beta) < 1+p$  and  $0 < C < C_*$ .

First, we establish the following rough estimation:

$$\begin{aligned} \left[ C^{1-\beta}(-x)_+^{\frac{(1+p)(1-\beta)}{mp-\beta}} - b(1-\beta)(1-(C/C_*)^{mp-\beta})t \right]_+^{\frac{1}{1-\beta}} \leq \\ \leq u(x,t) \leq C(-x)_+^{\frac{1+p}{mp-\beta}}, \text{ for } x \in \mathbb{R}, 0 \leq t < +\infty. \end{aligned} \quad (2.91)$$

To prove the left-hand side we consider the function,  $g$ , as in the case when  $p(m+\beta) > 1+p$  with  $\gamma = (C/C_*)^{mp-\beta}$ . As before, we then derive (2.89a), and since

$$S_t \leq 0, \text{ in } M,$$

we have  $S \leq 0$  in  $M$ . Hence, (2.90) is valid with  $\leq$  in (2.90a). As before, from Lemma 2.3.1, the left-hand side of (2.91) follows. To prove the right-hand side of (2.91) it is enough to observe that

$$Lu_0 = bu_0^\beta(1-(C/C_*)^{mp-\beta}) \geq 0, \text{ for } x \in \mathbb{R}, t \geq 0.$$

Having (2.91), we can now establish a more accurate estimation, (2.22). Consider a

function

$$g(x, t) = C_0 \left( -\zeta_0 t^{\frac{mp-\beta}{(1+p)(1-\beta)}} - x \right)_+^{\frac{1+p}{mp-\beta}}, \text{ in } G_\ell,$$

$$G_\ell := \left\{ (x, t) : \zeta(t) = -\ell t^{\frac{mp-\beta}{(1+p)(1-\beta)}} < x < +\infty, 0 < t < +\infty \right\},$$

where,  $C_0, \zeta_0 > 0$  and  $\ell > \zeta_0$ . Calculating  $Lg$  in

$$G_\ell^+ := \left\{ (x, t) : \zeta(t) < x < -\zeta_0 t^{\frac{mp-\beta}{(1+p)(1-\beta)}}, 0 < t < +\infty \right\},$$

we have

$$\begin{aligned} Lg = bg^\beta S, \quad S = 1 - (C_0/C_*)^{mp-\beta} - (b(1-\beta))^{-1} C_0^{1-\beta} \zeta_0 t^{\frac{p(m+\beta)-(1+p)}{(1+p)(1-\beta)}} \times \\ \times \left( -\zeta_0 t^{\frac{mp-\beta}{(1+p)(1-\beta)}} - x \right)^{\frac{1+p-p(m+\beta)}{mp-\beta}}. \end{aligned} \quad (2.92)$$

By choosing  $C_0 = C_*$ , we have

$$Lg \leq 0, \text{ in } G_\ell^+; \quad Lg = 0, \text{ in } G_\ell \setminus \bar{G}_\ell^+. \quad (2.93)$$

To obtain a lower estimation we choose  $\zeta_0 = \zeta_3$  and  $\ell = \ell_0$  (see Appendix 6.1). Using (2.91), we have

$$\begin{aligned} g|_{x=\zeta(t)} = t^{\frac{1}{1-\beta}} C_* (\ell_0 - \zeta_3)^{\frac{1+p}{mp-\beta}} = \left( b(1-\beta)\theta_* t \right)^{\frac{1}{1-\beta}} = t^{\frac{1}{1-\beta}} \times \\ \times \left[ C^{1-\beta} \ell_0^{\frac{(1+p)(1-\beta)}{mp-\beta}} - b(1-\beta) \left( 1 - (C/C_*)^{mp-\beta} \right) \right]^{\frac{1}{1-\beta}} \leq u(\zeta(t), t), \quad t \geq 0, \end{aligned} \quad (2.94a)$$

$$g(x, 0) = u(x, 0) = 0, \quad 0 \leq x \leq x_0, \quad (2.94b)$$

$$g(x_0, t) = u(x_0, t) = 0, \quad t \geq 0, \quad (2.94c)$$



where  $x_0 > 0$  is an arbitrary fixed number. By using (2.93) and (2.94), we can apply Lemma 2.3.1 in

$$G'_{\ell_0} := G_{\ell_0} \cap \{x < x_0\}.$$

Since  $x_0 > 0$  is arbitrary, the left inequality in (2.22) follows. Since

$$S_x \geq 0, \text{ for } \zeta(t) < x < -\zeta_0 t^{\frac{mp-\beta}{(1+p)(1-\beta)}}, t > 0,$$

from (2.92) it follows that

$$S \geq S|_{x=\zeta(t)} = 1 - (C_0/C_*)^{mp-\beta} - (b(1-\beta))^{-1} C_0^{1-\beta} \zeta_0 (\ell - \zeta_0)^{\frac{1+p-p(m+\beta)}{mp-\beta}}.$$

By choosing  $C_0 = C_3$ ,  $\zeta_0 = \zeta_4$ ,  $\ell = \ell_1$  (see Appendix 6.1), we have

$$S|_{x=\zeta(t)} = 0,$$

$$Lg \geq 0 \text{ in } G_{\ell_1}^+, Lg = 0 \text{ in } G_{\ell_1} \setminus \bar{G}_{\ell_1}^+,$$

$$u(\zeta(t), t) \leq t^{\frac{1}{1-\beta}} C_1^{\frac{1+p}{mp-\beta}} = t^{\frac{1}{1-\beta}} C_3 (\ell_1 - \zeta_4)^{\frac{1+p}{mp-\beta}} = g(\zeta(t), t), t \geq 0,$$

and, for arbitrary  $x_0 > 0$ , (2.94b) and (2.94c) are valid. By applying Lemma 2.3.1 in  $G'_{\ell_1}$ , due to the arbitrariness of  $x_0 > 0$ , we derive the right-hand side of (2.22). From (2.19), (2.21), and (2.22) it follows that

$$\zeta_1 t^{\frac{mp-\beta}{(1+p)(1-\beta)}} \leq \eta(t) \leq \zeta_2 t^{\frac{mp-\beta}{(1+p)(1-\beta)}}, 0 \leq t < +\infty,$$

where the constants  $\zeta_1$  and  $\zeta_2$  are chosen according to relevant estimations for  $u$ . From (2.16) and the respective estimations (2.19), (2.21), and (2.22), the estimation (2.20) fol-

lows. If  $u_0$  satisfies (1.9) with  $\alpha = (1+p)/(mp-\beta)$  and with  $C \neq C_*$ , then the asymptotic formulae (2.4) and (2.5) may be proved as the similar estimations (2.1) and (2.3) were in Lemma 2.3.4.  $\square$

### 2.4.3 Proof of Theorem 2.1.3

*Proof of Theorem 2.1.3.* For any  $\varepsilon > 0$  from (1.9), (2.37) follows. Consider a function

$$g_\varepsilon(x, t) = \left[ (C + \varepsilon)^{1-\beta} (-x)_+^{\alpha(1-\beta)} - b(1-\beta)(1-\varepsilon)t \right]_+^{1/(1-\beta)}. \quad (2.95)$$

We estimate  $Lg$  in

$$M_1 := \{(x, t) : x_\varepsilon < x < \eta_\ell(t), 0 < t \leq \delta_1\},$$

$$\eta_\ell(t) = -\ell t^{1/(\alpha(1-\beta))}, \quad \ell(\varepsilon) = (C + \varepsilon)^{-1/\alpha} [b(1-\beta)(1-\varepsilon)]^{1/\alpha(1-\beta)},$$

where  $\delta_1 > 0$  is chosen such that  $\eta_{\ell(\varepsilon)}(\delta_1) = x_\varepsilon$ . We have

$$Lg_\varepsilon = b g_\varepsilon^\beta \{\varepsilon + S\},$$

$$S = -b^{-1}(\alpha m)^p (C + \varepsilon)^{mp-\beta} (-x)_+^{\alpha(mp-\beta)-(1+p)} \left\{ g|x|^{-\alpha} / (C + \varepsilon) \right\}^{p(m+\beta)-(1+p)} S_1,$$

$$S_1 = \left\{ \alpha p(m+\beta-1) + p(\alpha(1-\beta)-1) \left[ g|x|^{-\alpha} / (C + \varepsilon) \right]^{1-\beta} \right\}.$$

If  $p(m+\beta) \geq 1+p$ , we can choose  $x_\varepsilon < 0$  such that

$$|S| < \frac{\varepsilon}{2}, \quad \text{in } M_1.$$

Thus we have

$$Lg_\varepsilon > b(\varepsilon/2)g_\varepsilon^\beta \text{ (respectively, } Lg_{-\varepsilon} < -b(\varepsilon/2)g_{-\varepsilon}^\beta), \text{ in } M_1, \quad (2.96)$$

$$Lg_{\pm\varepsilon} = 0, \text{ for } x > \eta_{\ell(\pm\varepsilon)}(t), \text{ } 0 < t \leq \delta_1, \quad (2.97)$$

$$g_\varepsilon(x, 0) \geq u_0(x) \text{ (respectively, } g_{-\varepsilon}(x, 0) \leq u_0(x)), \text{ } x \geq x_\varepsilon. \quad (2.98)$$

Since  $u$  and  $g$  are continuous functions,  $\delta = \delta(\varepsilon) \in (0, \delta_1]$ , may be chosen such that

$$g_\varepsilon(x_\varepsilon, t) \geq u(x_\varepsilon, t) \text{ (respectively, } g_{-\varepsilon}(x_\varepsilon, t) \leq u(x_\varepsilon, t)), \text{ } 0 \leq t \leq \delta.$$

From Lemma 2.3.1 it follows that

$$g_{-\varepsilon} \leq u \leq g_\varepsilon, \text{ } x \geq x_\varepsilon, \text{ } 0 \leq t \leq \delta, \quad (2.99a)$$

$$\eta_{\ell(-\varepsilon)}(t) \leq \eta(t) \leq \eta_{\ell(\varepsilon)}, \text{ } 0 \leq t \leq \delta, \quad (2.99b)$$

which imply (2.6) and (2.7). Let  $p(m+\beta) < 1+p$ . In this case the left-hand side of (2.99) may be proved similarly. Moreover, we can replace  $1+\varepsilon$  with  $1$  in  $g_{-\varepsilon}$  and  $\eta_{\ell(-\varepsilon)}$ . For a sharp upper estimation, consider a function

$$g(x, t) = C_6 \left( -\zeta_5 t^{\frac{1}{\alpha(1-\beta)}} - x \right)_+^\alpha, \text{ in } G_{\ell, \delta},$$

$$G_{\ell, \delta} := \{(x, t) : \eta_\ell(t) < x < +\infty, \text{ } 0 < t < \delta\},$$

where  $\ell \in (\ell_*, +\infty)$ ,  $C_6$  and  $\zeta_5$  are given in Appendix 6.1. From (2.7) it follows that for all  $\ell > \ell_*$  and for all sufficiently small  $\varepsilon > 0$ , there exists a  $\delta = \delta(\varepsilon, \ell) > 0$  such that

$$u(\eta_\ell(t), t) \leq t^{\frac{1}{1-\beta}} [C^{1-\beta} \ell^{\alpha(1-\beta)} - b(1-\beta)(1-\varepsilon)]^{\frac{1}{1-\beta}}, \text{ } 0 \leq t \leq \delta. \quad (2.100)$$

Calculating  $Lg$  in

$$G_{\ell, \delta}^+ = \left\{ (x, t) : \eta_\ell(t) < x < -\zeta_5 t^{\frac{1}{\alpha(1-\beta)}}, 0 < t < \delta \right\},$$

we derive

$$\begin{aligned} Lg &= bg^\beta S, \quad S = 1 - (b(1-\beta))^{-1} \zeta_5 C_6^{1/\alpha} \left\{ g t^{1/(\beta-1)} \right\}^{(\alpha(1-\beta)-1)/\alpha} \\ &\quad - \frac{p}{b} (\alpha m)^p (\alpha m - 1) C_6^{(1+p)/\alpha} g^{mp-\beta - ((1+p)/\alpha)}. \end{aligned}$$

Since

$$\begin{aligned} S_x &\geq 0, \quad \text{in } G_{\ell, \delta}^+, \\ S &\geq S \Big|_{x=\eta_\ell(t)} = 1 - (b(1-\beta))^{-1} \zeta_5 C_6^{1-\beta} (\ell - \zeta_5)^{\alpha(1-\beta)-1} - \\ &\quad - t^{\frac{\alpha(mp-\beta)-(1+p)}{\alpha(1-\beta)}} b^{-1} p (\alpha m)^p (\alpha m - 1) C_6^{mp-\beta} (\ell - \zeta_5)^{\alpha(mp-\beta)-(1+p)}, \end{aligned}$$

we have

$$S \geq \varepsilon - t^{\frac{\alpha(mp-\beta)-(1+p)}{\alpha(1-\beta)}} b^{-1} p (\alpha m)^p (\alpha m - 1) C_6^{mp-\beta} (\ell - \zeta_5)^{\alpha(mp-\beta)-(1+p)}, \quad \text{in } G_{\ell, \delta}^+.$$

By choosing  $\delta = \delta(\varepsilon) > 0$  sufficiently small we have

$$Lg \geq b(\varepsilon/2)g^\beta, \quad \text{in } G_{\ell, \delta}^+. \quad (2.101)$$

By applying (2.100) and Lemma 2.3.1 in  $G'_{\ell,\delta} = G_{\ell,\delta} \cap \{x < x_0\}$  we have

$$Lg = 0, \text{ in } G'_{\ell,\delta} \setminus \bar{G}_{\ell,\delta}^+, \quad (2.102a)$$

$$\begin{aligned} u(\eta_\ell(t), t) &\leq t^{\frac{1}{1-\beta}} [C^{1-\beta} \ell^{\alpha(1-\beta)} - b(1-\beta)(1-\varepsilon)]^{\frac{1}{1-\beta}} = \\ &= C_6(\ell - \zeta_5)^\alpha t^{\frac{1}{\alpha(1-\beta)}} = g(\eta_\ell(t), t), \quad 0 \leq t \leq \delta, \end{aligned} \quad (2.102b)$$

$$u(x_0, t) = g(x_0, t) = 0, \quad 0 \leq t \leq \delta, \quad (2.102c)$$

$$u(x, 0) = g(x, 0) = 0, \quad 0 \leq x \leq x_0. \quad (2.102d)$$

Since  $x_0 > 0$  is arbitrary, from (2.102) and Lemma 2.3.1, it follows that for all  $\ell > \ell_*$  and  $\varepsilon > 0$ , there exists  $\delta = \delta(\varepsilon, \ell) > 0$  such that

$$u(x, t) \leq C_6 \left( -\zeta_5 t^{\frac{1}{\alpha(1-\beta)}} - x \right)_+^\alpha, \text{ in } \bar{G}_{\ell,\delta}. \quad (2.103)$$

In view of (2.7) (which is valid along  $x = \eta_\ell(t)$ ),  $\delta$  may be chosen so small that

$$-\ell t^{1/\alpha(1-\beta)} \leq \eta(t) \leq -\zeta_5 t^{1/\alpha(1-\beta)}, \quad 0 \leq t \leq \delta. \quad (2.104)$$

Since  $\ell > \ell_*$  and  $\varepsilon > 0$  are arbitrary, (2.6) follows from (2.104).  $\square$

The proofs of Theorem 2.1.4 and estimations (2.23)-(2.32) are almost identical to the similar proofs given in [23].

## Chapter 3

# Traveling-Wave Solutions to the Nonlinear Double Degenerate Reaction-Diffusion Equation

### 3.1 Introduction and the Main Result

In this section we consider equation (1.7)

$$u_t = \left( |(u^m)_x|^{p-1} (u^m)_x \right)_x - bu^\beta, \quad x \in \mathbb{R}, t > 0,$$

with  $mp > 1$ ,  $(m, p > 0)$ ,  $0 < \beta < 1$ , and  $b > 0$ . We present asymptotic results for the finite traveling-wave solutions to (1.7):  $u(x, t) = \varphi(kt - x)$ , where  $\varphi(y) \geq 0$ ,  $\varphi \not\equiv 0$ , and  $\varphi(y) \rightarrow 0^+$  as  $y \rightarrow -\infty$ ;  $\varphi(y) = 0$  for  $y \leq y_0$ , for some  $y_0 \in \mathbb{R}$ . By translation we select  $y_0 = 0$ . Finite traveling-wave solutions are weak solutions to (1.7), (1.8) in sense of Definition 1.2.1.

There exists a finite traveling-wave solution to equation (1.7) if we can find a func-

tion  $\varphi \in \mathbb{R}^+$  that satisfies the following initial-value problem (IVP)

$$\begin{cases} \left( |(\varphi^m)'|^{p-1} (\varphi^m)' \right)' - k\varphi' - b\varphi^\beta = 0, \\ \varphi(0) = (\varphi^m)'(0) = 0, \end{cases} \quad (3.1)$$

we extend  $\varphi(y)$  by 0 on for all  $y < 0$ . Note that all derivatives are understood in the weak sense.

The following is the main result of this Chapter.

**Theorem 3.1.1.** *There exists a finite traveling-wave solution,  $u(x, t) = \varphi(kt - x)$ , to equation (1.7), with  $\varphi(0) = 0$  if  $k \neq 0$ . Further we have,*

1.  $\lim_{y \rightarrow 0^+} y^{-\frac{1+p}{mp-\beta}} \varphi(y) = C_* = \left[ \frac{b(mp-\beta)^{1+p}}{(m(1+p))^p p(m+\beta)} \right]^{\frac{1}{mp-\beta}}$ , if  $p(m+\beta) < 1+p$ ;
2.  $\lim_{y \rightarrow +\infty} y^{-\frac{1+p}{mp-\beta}} \varphi(y) = C_*$ , if  $p(m+\beta) > 1+p$ ;
3.  $\lim_{y \rightarrow +\infty} y^{-\frac{p}{mp-1}} \varphi(y) = \left( \frac{mp-1}{mp} \right)^{\frac{p}{mp-1}} k^{\frac{1}{mp-1}}$ , if  $k > 0$ ,  $p(m+\beta) < 1+p$ ;
4.  $\lim_{y \rightarrow 0^+} y^{-\frac{p}{mp-1}} \varphi(y) = \left( \frac{mp-1}{mp} \right)^{\frac{p}{mp-1}} k^{\frac{1}{mp-1}}$ , if  $k > 0$ ,  $p(m+\beta) > 1+p$ ;
5.  $\lim_{y \rightarrow +\infty} y^{-\frac{1}{1-\beta}} \varphi(y) = \left( (1-\beta) \left( -\frac{b}{k} \right) \right)^{\frac{1}{1-\beta}}$ , if  $k < 0$ ,  $p(m+\beta) < 1+p$ ;
6.  $\lim_{y \rightarrow 0^+} y^{-\frac{1}{1-\beta}} \varphi(y) = \left( (1-\beta) \left( -\frac{b}{k} \right) \right)^{\frac{1}{1-\beta}}$ , if  $k < 0$ ,  $p(m+\beta) > 1+p$ .

The existence of traveling-wave solutions with interfaces for the nonlinear diffusion equation ((1.7) with  $p = 1$ ) is considered in [65] and for the parabolic  $p$ -Laplacian equation ((1.7) with  $m = 1$ ) in [80].

The organization of this section is as follows: In Section 3.1.1 we formulate and prove some preliminary results which are necessary for the proof of main result and in

Section 3.2 we prove the main result, Theorem 3.1.1.

### 3.1.1 Preliminary Results: Traveling-Wave Solutions and Phase-Space Analysis

In this section we'll apply phase-space analysis to find finite traveling-wave solutions for (1.7). We aim to analyze the phase portrait for problem (3.1). Consider the following lemma.

**Lemma 3.1.2.** *If the function  $\varphi(y)$  is a positive solution to problem (3.1), then  $\varphi(y)$  is increasing for all  $y > 0$ .*

The proof of Lemma 3.1.2 follows as in the proof of the analogous result for the  $p$ -Laplacian equation in [80].

There exists a unique  $\varphi(y) > 0$  that satisfies the IVP (3.1) for  $y > 0$ . To prove this, we introduce the following change of variable

$$\Theta = \varphi \text{ and } \Upsilon = ((\varphi^m)')^p,$$

it follows that

$$\Theta' = \frac{1}{m} \Theta^{1-m} \Upsilon^{\frac{1}{p}} \text{ and } \Upsilon' = b \Theta^\beta + \frac{k}{m} \Theta^{1-m} \Upsilon^{\frac{1}{p}}.$$

$(\Theta, \Upsilon)$  starts from  $(0,0)$  at  $y = 0$ , exists for any  $y \in \mathbb{R}^+$ , and are contained in the first quadrant:  $Q_1 = \{(\Theta, \Upsilon) : \Theta, \Upsilon > 0\}$  for  $y > 0$ . There exists a unique solution, or trajectory,



$\Upsilon(\Theta)$ . To prove this, we divide  $\Upsilon'$  by  $\Theta'$  and derive the following problem

$$\begin{cases} \frac{d\Upsilon}{d\Theta} = f(\Theta, \Upsilon) = k + bm\Theta^{m+\beta-1}\Upsilon^{-\frac{1}{p}}, \\ \Upsilon(0) = 0. \end{cases} \quad (3.2)$$

As done in [80], we find the nonzero solutions or trajectories,  $\Upsilon(\Theta)$ , to (3.2), in two steps. First, we prove the global existence of the solution of the following perturbed IVP

$$\begin{cases} \frac{d\Upsilon}{d\Theta} = f(\Theta, \Upsilon) = k + bm\Theta^{m+\beta-1}\Upsilon^{-\frac{1}{p}}, \\ \Upsilon(0) = \varepsilon, \varepsilon > 0. \end{cases} \quad (3.3)$$

The fact that the function  $f(\Theta, \Upsilon)$  is locally Lipschitz continuous in  $\mathbb{R}^+ \times (\varepsilon, +\infty)$  implies that there exists a unique local solution  $\Upsilon_\varepsilon$  to problem (3.3). For both  $k > 0$  and for  $k < 0$  with  $p(m+\beta) > 1+p$ , the proof of the existence of a global solution to (3.3) follows as in the proof of the existence of a global solution to the analogous IVP for the  $p$ -Laplacian equation in [80].

Let  $k < 0$  with  $p(m+\beta) < 1+p$ . The difference from the previous case is that

$$\tilde{C} : (0, +\infty) \rightarrow (+\infty, 0),$$

see Figure 3.1b. Since

$$\frac{d\Upsilon_\varepsilon}{d\Theta} > 0 \text{ if } \Upsilon_\varepsilon < \left( -\frac{k}{bm}\Theta^{1-m-\beta} \right)^{-p},$$

i.e.,  $\Upsilon_\varepsilon$  is increasing to the left of  $\tilde{C}$ . Then  $\Upsilon_\varepsilon$  must cross  $\tilde{C}$  with horizontal tangent, after that  $\Upsilon_\varepsilon$  will be strictly decreasing. It follows that  $\Upsilon_\varepsilon$  is a global solution to (3.3) if  $k < 0$ .

We now prove the global existence of solution to the perturbed problem

$$\begin{cases} \frac{d\Upsilon}{d\Theta} = f(\Theta, \Upsilon) = k + bm\Theta^{m+\beta-1}\Upsilon^{-\frac{1}{p}}, \\ \Upsilon(\varepsilon) = 0, \varepsilon > 0. \end{cases} \quad (3.4)$$

To do this, we consider the following Cauchy problem for the inverse function of  $\Upsilon$ , denoted as  $v$

$$\begin{cases} \frac{dv}{dt} = g(v, t) = \frac{1}{f(v, t)} = \frac{t^{\frac{1}{p}}}{kt^{\frac{1}{p}} + bmv^{m+\beta-1}}, \\ v(0) = \varepsilon, \varepsilon > 0. \end{cases} \quad (3.5)$$

Since the right hand side of (3.5) is Lipschitz continuous, there exists a local solution,  $v_\varepsilon$ , to the CP (3.5). For  $k > 0$  and for  $k < 0$  with  $p(m+\beta) > 1+p$ , as for (3.3), the proof of the existence of a global solution to (3.4) follows as in the proof of the existence of a global solution to the analogous IVP for the  $p$ -Laplacian equation in [80].

Let  $k < 0$  with  $p(m+\beta) < 1+p$ . As before, we define the curve where  $f(v, t) = 0$  by  $\tilde{C}$ . We denote the region to the left of  $\tilde{C}$  as  $Q_l = \{(v, t) : f(v, t) > 0\}$  and to the region to the right of  $\tilde{C}$  as  $Q_r = \{(v, t) : f(v, t) < 0\}$ , see Figure 3.1d. Since  $v$  is increasing in  $Q_l$  it must cross  $\tilde{C}$  with vertical tangent, however, this is impossible. Let  $t_\varepsilon$  be such that  $v(t_\varepsilon) = M_\varepsilon \in \tilde{C}$ . Consider the function  $w$  such that

$$w : [\varepsilon, M_\varepsilon] \rightarrow [0, t_\varepsilon].$$

Then  $w$  is the inverse function of  $v$  in  $[0, t_\varepsilon]$  and so solves the following problem

$$\begin{cases} \frac{dw}{dt} = k + bmw^{m+\beta-1}t^{-\frac{1}{p}} = f(w, t), \\ w(\varepsilon) = 0, w(M_\varepsilon) = t_\varepsilon, \varepsilon > 0. \end{cases} \quad (3.6)$$

Let  $\widehat{C}$  denote the curve where  $f(w, t) = 0$ . So  $w$  enters the region to the right of  $\widehat{C}$  with horizontal tangent and since if  $t > M_\varepsilon$ , then  $w(t)$  is decreasing, we have that  $w$  cannot cross  $\widehat{C}$  again since it must cross with horizontal tangent, which is a contradiction. It follows that the solution,  $w$ , to problem (3.6) is global and so there exists a global solution to problem (3.4) if  $k < 0$ .

**Lemma 3.1.3.** *The problem (3.2) has a unique global solution.*

The proof of Lemma 3.1.3 follows as in the proof of existence and uniqueness of solution for the analogous problem for the  $p$ -Laplacian equation in [80].




Figure 3.1: Trajectories  $\Upsilon(\Theta)$  and  $v(t)$

Let  $\Upsilon = ((\varphi^m)')^p$  be a solution of the problem (3.2). For the problem

$$\frac{d\varphi}{dy} = \frac{1}{m}(\varphi(y))^{1-m}\Upsilon^{\frac{1}{p}}(\varphi(y)), \varphi(0) = 0, \quad (3.7)$$

there exists a unique maximal solution defined on  $(-\infty, \ell)$  such that

$$\lim_{y \rightarrow \ell^-} \varphi(y) = +\infty.$$

By (3.7) we have that  $(\varphi^m)'(0) = \Upsilon^{\frac{1}{p}}(0) = 0$ , so we can continue  $\varphi$  by zero on  $(-\infty, 0)$ .

On the other side,  $\varphi$  is strictly increasing, and

$$\lim_{y \rightarrow \ell^-} \varphi(y) = +\infty,$$

if  $\ell$  is finite. By (3.7) and the boundedness of  $\Upsilon^{-\frac{1}{p}}$ , the above limit also holds if  $\ell = +\infty$ .

The solution of (3.7) defined on  $(-\infty, \ell)$  satisfies the following problem

$$\begin{cases} (|(\varphi^m)'|^{p-1}(\varphi^m)')' - k\varphi' - b\varphi^\beta = 0, \text{ on } (-\infty, \ell), \\ \varphi(0) = (\varphi^m)'(0) = 0. \end{cases} \quad (3.8)$$

We now prove that the solution to problem (3.8) is global. To proceed, we require the following lemma.

**Lemma 3.1.4.** *Let  $\Upsilon$  be a solution of the problem (3.2). Then we have*

1.  $\Upsilon(\Theta) \sim \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}} \Theta^{\frac{p(m+\beta)}{1+p}}$ , as  $\Theta \rightarrow 0^+$  if  $p(m+\beta) < 1+p$ .
2.  $\Upsilon(\Theta) \sim \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}} \Theta^{\frac{p(m+\beta)}{1+p}}$ , as  $\Theta \rightarrow +\infty$  if  $p(m+\beta) > 1+p$ .
3.  $\Upsilon(\Theta) \sim k\Theta$ , as  $\Theta \rightarrow +\infty$  if  $k > 0$ ,  $p(m+\beta) < 1+p$ .
4.  $\Upsilon(\Theta) \sim k\Theta$ , as  $\Theta \rightarrow 0^+$  if  $k > 0$ ,  $p(m+\beta) > 1+p$ .
5.  $\Upsilon(\Theta) \sim \left(-\frac{k}{bm}\right)^{-p} \Theta^{p(m+\beta-1)}$ , as  $\Theta \rightarrow +\infty$  if  $k < 0$ ,  $p(m+\beta) < 1+p$ .
6.  $\Upsilon(\Theta) \sim \left(-\frac{k}{bm}\right)^{-p} \Theta^{p(m+\beta-1)}$ , as  $\Theta \rightarrow 0^+$  if  $k < 0$ ,  $p(m+\beta) > 1+p$ .

*Proof of Lemma 3.1.4.* We begin by proving formulas 1. and 2. from Lemma 3.1.4. We apply nonlinear scaling as follows: we choose  $\Upsilon_l(\Theta) = l\Upsilon(l^\gamma\Theta)$ , with  $l > 0$  and  $\gamma$  to be determined.

$$\Upsilon_l(\Theta) = l^\gamma \Upsilon(l^\gamma \Theta) \iff \Upsilon(\Theta) = l^{-1} \Upsilon_l(l^{-\gamma} \Theta).$$

We set  $Z = l^\gamma \Theta$ . It follows from (3.2) that

$$\begin{aligned} \frac{d\Upsilon_l}{d\Theta} &= l^{1+\gamma} \frac{d\Upsilon}{dZ} = l^{1+\gamma} \left( k + bmZ^{m+\beta-1} \Upsilon^{-\frac{1}{p}} \right) \\ &= kl^{1+\gamma} + bml^{1+\gamma} l^{\gamma(m+\beta-1)} l^{\frac{1}{p}} \Theta^{m+\beta-1} \Upsilon_l^{-\frac{1}{p}}. \end{aligned} \quad (3.9)$$

We choose  $\gamma$  such that

$$1 + \gamma + \gamma(m + \beta - 1) + \frac{1}{p} = 0 \implies \gamma = -\frac{1+p}{p(m+\beta)}.$$

So we have that

$$\frac{d\Upsilon_l}{d\Theta} = kl^{\frac{p(m+\beta)-(1+p)}{p(m+\beta)}} + bm\Theta^{m+\beta-1} \Upsilon_l^{-\frac{1}{p}}. \quad (3.10)$$

From our previous results we that there exists a unique solution to (3.10). To prove formula 1., since  $p(m + \beta) < 1 + p$ , we set

$$\lim_{l \rightarrow +\infty} \Upsilon_l(\Theta) = \tilde{\Upsilon}(\Theta),$$

where  $\tilde{\Upsilon}(\Theta)$  solves

$$\begin{cases} \frac{d\Upsilon}{d\Theta} = bm\Theta^{m+\beta-1} \Upsilon^{-\frac{1}{p}}, \\ \Upsilon(0) = 0. \end{cases} \quad (3.11)$$

The existence of the above limit follows from a similar argument used to prove an analogous limit in the proof of formula 3. The ODE in (3.11) is separable. Separating variables and integrating we have that

$$\tilde{\Upsilon}(\Theta) = \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}} \Theta^{\frac{p(m+\beta)}{1+p}}. \quad (3.12)$$

Recall that  $Z = l^\gamma \Theta \implies \Theta = l^{-\gamma} Z$ . So we have that

$$\Theta^{\frac{p(m+\beta)}{1+p}} = l^{-\frac{\gamma p(m+\beta)}{1+p}} Z^{\frac{p(m+\beta)}{1+p}} = l Z^{\frac{p(m+\beta)}{1+p}}.$$

It follows that

$$\begin{aligned} \lim_{l \rightarrow +\infty} \Upsilon_l(\Theta) &= \lim_{l \rightarrow +\infty} l \Upsilon(Z) = \widetilde{\Upsilon}(\Theta) \\ &= \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}} l Z^{\frac{p(m+\beta)}{1+p}} \\ \implies \lim_{Z \rightarrow 0^+} \frac{\Upsilon(Z)}{Z^{\frac{p(m+\beta)}{1+p}}} &= \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}}. \end{aligned}$$

Therefore,

$$\Upsilon(\Theta) \sim \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}} \Theta^{\frac{p(m+\beta)}{1+p}}, \text{ as } \Theta \rightarrow 0^+.$$

Note that formula 2, where  $p(m+\beta) > 1+p$ , follows from the same procedure by setting

$$\lim_{l \rightarrow 0^+} \Upsilon_l(\Theta) = \widetilde{\Upsilon}(\Theta).$$

To prove formulas 3. and 4. we let  $k > 0$  and proceed as in the proof of formulas 1. and 2. We choose the same scale as follows

$$\Upsilon_l(\Theta) = l \Upsilon(l^\gamma \Theta) \iff \Upsilon(\Theta) = l^{-1} \Upsilon_l(l^{-\gamma} \Theta).$$

We set  $Z = l^\gamma \Theta$ . It follows from (3.2) that

$$\begin{aligned} \frac{d\Upsilon_l}{d\Theta} &= l^{1+\gamma} \frac{d\Upsilon}{dZ} = l^{1+\gamma} \left( k + bmZ^{m+\beta-1} \Upsilon^{-\frac{1}{p}} \right) \\ &= kl^{1+\gamma} + bml^{1+\gamma} l^{\gamma(m+\beta-1)} l^{\frac{1}{p}} \Theta^{m+\beta-1} \Upsilon_l^{-\frac{1}{p}}. \end{aligned} \tag{3.13}$$

Now, we choose  $\gamma$  such that

$$1 + \gamma = 0 \implies \gamma = -1.$$

So we have that

$$\frac{d\Upsilon_l}{d\Theta} = k + bml^{\frac{1+p-p(m+\beta)}{p}} \Theta^{m+\beta-1} \Upsilon_l^{-\frac{1}{p}}. \quad (3.14)$$

From our previous results we that there exists a unique solution to (3.14). To prove formula 3., since  $p(m+\beta) < 1+p$ , we set

$$\lim_{l \rightarrow 0^+} \Upsilon_l(\Theta) = \widetilde{\Upsilon}(\Theta).$$

To prove the existence of this limit, let  $0 \leq \Gamma < \Delta < +\infty$ . We show

- $\{\Upsilon_l\}$  is uniformly bounded, i.e.,  $|\Upsilon_l(\Theta)| \leq C$ , for all  $\Theta \in [\Gamma, \Delta]$  and  $l$ , where  $C$  is independent of  $l$ .
- $\{\Upsilon_l\}$  is equicontinuous, i.e., for any  $\varepsilon > 0$ , there exists  $\delta = \delta_\varepsilon > 0$  such that for all  $\Theta, \Theta_0 \in [\Gamma, \Delta]$  we have

$$|\Theta - \Theta_0| < \delta \implies |\Upsilon_l(\Theta) - \Upsilon_l(\Theta_0)| < \varepsilon, \forall l.$$

First we prove that  $\{\Upsilon_l\}$  is uniformly bounded. Since we want to pass  $l$  to zero, we fix  $l \in (0, 1]$ . So we have that

$$\frac{d\Upsilon_l}{d\Theta} = k + bml^{\frac{1+p-p(m+\beta)}{p}} \Theta^{m+\beta-1} \Upsilon_l^{-\frac{1}{p}} \leq k + bm\Theta^{m+\beta-1} \Upsilon_1^{-\frac{1}{p}} = \frac{d\Upsilon_1}{d\Theta}.$$

Choosing  $\Gamma = 0$  we have that  $\Upsilon_l(0) = \Upsilon_1(0) = 0$ , so by applying the comparison theorem



we have

$$0 \leq \Upsilon_l(\Theta) \leq \Upsilon_1(\Theta), \forall \Theta \in [0, \Delta], \forall l \in (0, 1].$$

It remains to show that  $\frac{d\Upsilon_l}{d\Theta}$  is uniformly bounded. Let  $\Theta \in [\Gamma, \Delta]$ . Since  $k > 0$  we have that

$$\frac{d\Upsilon_l}{d\Theta} \geq k \implies \Upsilon_l(\Theta) \geq k\Theta \implies \Upsilon_l(\Gamma) \geq k\Gamma > 0 \implies \Upsilon_l^{-\frac{1}{p}}(\Gamma) \leq (k\Gamma)^{-\frac{1}{p}}.$$

So we have

$$\frac{d\Upsilon_l}{d\Theta} = k + bml^{\frac{1+p-p(m+\beta)}{p}} \Theta^{m+\beta-1} \Upsilon_l^{-\frac{1}{p}}(\Theta) \leq k + bm\Delta^{m+\beta-1} (k\Gamma)^{-\frac{1}{p}} < +\infty.$$

This holds for all  $l \in (0, 1]$ . Since  $\frac{d\Upsilon_l}{d\Theta}$  is uniformly bounded on  $[\Gamma, \Delta]$  it follows that  $\Upsilon_l(\Theta)$  is uniformly bounded on  $[\Gamma, \Delta]$ .

Now we need to show that  $\{\Upsilon_l\}$  is equicontinuous on  $[\Gamma, \Delta]$ . Let  $\Theta, \Theta_0 \in [\Gamma, \Delta]$ . We need to show that for any  $\varepsilon > 0$ , there exists  $\delta = \delta_\varepsilon > 0$  such that

$$|\Theta - \Theta_0| < \delta \implies |\Upsilon_l(\Theta) - \Upsilon_l(\Theta_0)| < \varepsilon, \forall l.$$

By Lagrange's mean value theorem, for all  $\theta \in [0, 1]$ , we have

$$|\Upsilon_l(\Theta) - \Upsilon_l(\Theta_0)| = \left| \frac{d\Upsilon_l(\Theta_0 + \theta(\Theta - \Theta_0))}{d\Theta} (\Theta - \Theta_0) \right| \leq C|\Theta - \Theta_0| < C\delta.$$

Choosing  $\delta = \frac{\varepsilon}{C}$  ensures that  $|\Upsilon_l(\Theta) - \Upsilon_l(\Theta_0)| < \varepsilon, \forall l$ . So  $\{\Upsilon_l\}$  is equicontinuous on  $[\Gamma, \Delta]$ .

Since  $\{\Upsilon_l\}$  is both uniformly bounded and equicontinuous on  $[\Gamma, \Delta]$ , and since  $[\Gamma, \Delta]$  is an arbitrary compact subset of  $[0, +\infty)$ , there exists  $\widetilde{\Upsilon}(\Theta)$  such that for some subsequence  $l'$  we have

$$\lim_{l' \rightarrow 0^+} \Upsilon_{l'}(\Theta) = \widetilde{\Upsilon}(\Theta), \forall \Theta > 0.$$

Where  $\tilde{\Upsilon}(\Theta)$  solves

$$\begin{cases} \frac{d\Upsilon}{d\Theta} = k, \Theta > 0, \\ \Upsilon(0) = 0. \end{cases} \quad (3.15)$$

So  $\tilde{\Upsilon}(\Theta) = k\Theta$ , and we have

$$\lim_{l \rightarrow 0^+} \Upsilon_l(\Theta) = \lim_{l \rightarrow 0^+} l\Upsilon(l^\gamma \Theta) = k\Theta, \Theta > 0.$$

Recall that  $Z = l^\gamma \Theta \implies \Theta = l^{-\gamma} Z$ . So we have that

$$\begin{aligned} \lim_{l \rightarrow 0^+} \Upsilon_l(\Theta) &= \lim_{l \rightarrow 0^+} l\Upsilon(Z) = \tilde{\Upsilon}(\Theta) = klZ \\ &\implies \lim_{Z \rightarrow +\infty} \frac{\Upsilon(Z)}{Z} = k. \end{aligned}$$

Therefore,

$$\Upsilon(\Theta) \sim k\Theta, \text{ as } \Theta \rightarrow +\infty.$$

Note that formula 4., where  $p(m + \beta) > 1 + p$ , follows from the same procedure by setting

$$\lim_{l \rightarrow +\infty} \Upsilon_l(\Theta) = \tilde{\Upsilon}(\Theta).$$

To prove formulas 5. and 6. we let  $k < 0$  and proceed as in the proof of the previous formulas. We choose the same scale as follows

$$\Upsilon_l(\Theta) = l\Upsilon(l^\gamma \Theta) \iff \Upsilon(\Theta) = l^{-1}\Upsilon_l(l^{-\gamma}\Theta).$$

We set  $Z = l^\gamma \Theta$ . It follows from (3.2) that

$$\begin{aligned} \frac{d\Upsilon_l}{d\Theta} &= l^{1+\gamma} \frac{d\Upsilon}{dZ} = l^{1+\gamma} \left( k + bmZ^{m+\beta-1} \Upsilon^{-\frac{1}{p}} \right) \\ &= kl^{1+\gamma} + bml^{1+\gamma} l^{\gamma(m+\beta-1)} l^{\frac{1}{p}} \Theta^{m+\beta-1} \Upsilon_l^{-\frac{1}{p}}. \end{aligned} \quad (3.16)$$

Now, we choose  $\gamma$  such that

$$1 + \gamma = 1 + \gamma + \gamma(m + \beta - 1) + \frac{1}{p} \implies \gamma = -\frac{1}{p(m + \beta - 1)}.$$

So we have that

$$l^{\frac{1+p-p(m+\beta)}{p(m+\beta-1)}} \frac{d\Upsilon_l}{d\Theta} = k + bm\Theta^{m+\beta-1} \Upsilon_l^{-\frac{1}{p}}. \quad (3.17)$$

From our previous results we that there exists a unique solution to (3.17). To prove formula 5., since  $p(m + \beta) < 1 + p$ , we set

$$\lim_{l \rightarrow 0^+} \Upsilon_l(\Theta) = \tilde{\Upsilon}(\Theta).$$

As before, we have to show that the above limit exists. In this case, it's enough to prove that  $\{\Upsilon_l\}$  is uniformly bounded on any compact interval,  $[\Gamma, \Delta]$ . From the equation we have that

$$k + bm\Theta^{m+\beta} \Upsilon_l^{-\frac{1}{p}} \geq 0 \implies 0 \leq \Upsilon_l(\Theta) \leq \left( -\frac{k}{bm} \right)^{-p} \Theta^{p(m+\beta-1)}, \Theta > 0.$$

It remains to show that  $\frac{d\Upsilon_l}{d\Theta}$  is uniformly bounded on  $[\Gamma, \Delta]$ . Consider

$$l^{\frac{1+p-p(m+\beta)}{p(m+\beta-1)}} \frac{d\Upsilon_l}{d\Theta} = k + bm\Theta^{m+\beta-1} \Upsilon_l^{-\frac{1}{p}} \implies \frac{d\Upsilon_l}{d\Theta} = l^{\frac{p(m+\beta)-(1+p)}{p(m+\beta-1)}} \left( k + bm\Theta^{m+\beta-1} \Upsilon_l^{-\frac{1}{p}} \right),$$

$$\begin{aligned}
(l+1)^{\frac{1+p-p(m+\beta)}{p(m+\beta-1)}} \frac{d\Upsilon_{l+1}}{d\Theta} &= k + bm\Theta^{m+\beta-1} \Upsilon_{l+1}^{-\frac{1}{p}} \\
\implies \frac{d\Upsilon_{l+1}}{d\Theta} &= (l+1)^{\frac{p(m+\beta)-(1+p)}{p(m+\beta-1)}} \left( k + bm\Theta^{m+\beta-1} \Upsilon_{l+1}^{-\frac{1}{p}} \right) \leq l^{\frac{p(m+\beta)-(1+p)}{p(m+\beta-1)}} \left( k + bm\Theta^{m+\beta-1} \Upsilon_{l+1}^{-\frac{1}{p}} \right).
\end{aligned}$$

Define  $Z(\Theta) := \Upsilon_{l+1}(\Theta) - \Upsilon_l(\Theta)$ . By mean value theorem, for all  $\theta \in [0, 1]$ , we have

$$\begin{aligned}
\frac{dZ}{d\Theta} &\leq l^{\frac{p(m+\beta)-(1+p)}{p(m+\beta-1)}} bm\Theta^{m+\beta-1} \left( \Upsilon_{l+1}^{-\frac{1}{p}} - \Upsilon_l^{-\frac{1}{p}} \right) = \\
&= -l^{\frac{p(m+\beta)-(1+p)}{p(m+\beta-1)}} \frac{bm}{p} \Theta^{m+\beta-1} (\Upsilon_l + \theta(\Upsilon_{l+1} - \Upsilon_l))^{-\frac{1+p}{p}} Z \\
\implies l^{\frac{1+p-p(m+\beta)}{p(m+\beta-1)}} \frac{dZ}{d\Theta} &\leq -\frac{bm}{p} \Theta^{m+\beta-1} (\Upsilon_l + \theta(\Upsilon_{l+1} - \Upsilon_l))^{-\frac{1+p}{p}} Z.
\end{aligned}$$

Since  $Z(0) = 0$ , it follows from the comparison theorem that  $\Upsilon_{l+1}(\Theta) \leq \Upsilon_l(\Theta)$ ,  $\Theta \in [\Gamma, \Delta]$ . Hence  $\{\Upsilon_l\}$  is a monotonically decreasing sequence as  $l \rightarrow 0^+$ , and since  $\Upsilon_l(\Theta) > 0$ , for all  $\Theta > 0$ , there exists  $\widetilde{\Upsilon}(\Theta)$  such that

$$\lim_{l \rightarrow 0^+} \Upsilon_l(\Theta) = \widetilde{\Upsilon}(\Theta).$$

Now, for any  $\nu \in C_0^\infty(\Gamma, \Delta)$ , we appeal to the integral identity

$$\int_{\Gamma}^{\Delta} l^{\frac{1+p-p(m+\beta)}{p(m+\beta-1)}} \Upsilon_l \nu' + (k + bm\Theta^{m+\beta-1} \Upsilon_l^{-\frac{1}{p}}) \nu d\Theta = 0.$$

Letting  $l \rightarrow 0^+$  we have

$$\int_{\Gamma}^{\Delta} (k + bm\Theta^{m+\beta-1} \widetilde{\Upsilon}^{-\frac{1}{p}}) \nu d\Theta = 0.$$

Since  $\nu$  is arbitrary we necessarily have that

$$k + bm\Theta^{m+\beta-1}\tilde{\Upsilon}^{-\frac{1}{p}} = 0.$$

Solving for  $\tilde{\Upsilon}$  we have that

$$\tilde{\Upsilon}(\Theta) = \left(-\frac{k}{bm}\right)^{-p} \Theta^{p(m+\beta-1)}. \quad (3.18)$$

Recall that  $Z = l^\gamma \Theta \implies \Theta = l^{-\gamma} Z$ . So we have that

$$\begin{aligned} \lim_{l \rightarrow 0^+} \Upsilon_l(\Theta) &= \lim_{l \rightarrow 0^+} l \Upsilon(Z) = \tilde{\Upsilon}(\Theta) = \left(-\frac{k}{bm}\right)^{-p} l Z^{p(m+\beta-1)} \\ &\implies \lim_{Z \rightarrow +\infty} \frac{\Upsilon(Z)}{Z^{p(m+\beta-1)}} = \left(-\frac{k}{bm}\right)^{-p}. \end{aligned}$$

Therefore,

$$\Upsilon(\Theta) \sim \left(-\frac{k}{bm}\right)^{-p} \Theta^{p(m+\beta-1)}, \text{ as } \Theta \rightarrow +\infty.$$

The proof of formula 6. follows from a similar argument. □

## 3.2 Proof of the Main Result

Using the results above we now prove Theorem 3.1.1.

*Proof of Theorem 3.1.1.* As long as  $\varphi(y) \neq 0$  ( $\Upsilon(\varphi(y)) \neq 0$ ), we write (3.7) as

$$m\varphi^{m-1}\Upsilon^{-\frac{1}{p}}(\varphi(y))d\varphi(y) = dy \quad (3.19)$$

We will prove formula 2. from Theorem 3.1.1, the proof of formula 1. is similar.

Since  $p(m+\beta) > 1 + p$ , from Lemma 3.1.4 we know that

$$\Upsilon(\Theta) \sim \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}} \Theta^{\frac{p(m+\beta)}{1+p}}, \text{ as } \Theta \rightarrow +\infty.$$

Hence, for any  $\varepsilon > 0$ , there exists an  $\Theta_\varepsilon \gg 1$  such that

$$\left( \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}} - \varepsilon \right) \Theta^{\frac{p(m+\beta)}{1+p}} \leq \Upsilon(\Theta) \leq \left( \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}} + \varepsilon \right) \Theta^{\frac{p(m+\beta)}{1+p}}, \Theta_\varepsilon \gg 1.$$

Raising both sides by the power  $-\frac{1}{p}$  then multiplying by  $m\Theta^{m-1}$  we have

$$\begin{aligned} m \left( \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}} + \varepsilon \right)^{-\frac{1}{p}} \Theta^{-\frac{m+\beta}{1+p}+m-1} &\leq m\Theta^{m-1}\Upsilon^{-\frac{1}{p}} \leq \\ m \left( \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}} - \varepsilon \right)^{-\frac{1}{p}} \Theta^{-\frac{m+\beta}{1+p}+m-1}, &\Theta_\varepsilon \gg 1. \end{aligned}$$

Integrating over  $(0, \varphi(y))$  where  $(0, y) \subset (0, \ell)$  we have

$$\begin{aligned} \frac{m(1+p)}{mp-\beta} \left( \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}} + \varepsilon \right)^{-\frac{1}{p}} (\varphi(y))^{\frac{mp-\beta}{1+p}} &\leq m \int_0^{\varphi(y)} \Theta^{m-1} \Upsilon^{-\frac{1}{p}}(\Theta) d\Theta \leq \\ \frac{m(1+p)}{mp-\beta} \left( \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}} - \varepsilon \right)^{-\frac{1}{p}} (\varphi(y))^{\frac{mp-\beta}{1+p}}. & \end{aligned}$$

Rewriting

$$\frac{m(1+p)}{mp-\beta} \left( \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}} + \varepsilon \right)^{-\frac{1}{p}} \leq \frac{m \int_0^{\varphi(y)} \Theta^{m-1} \Upsilon^{-\frac{1}{p}}(\Theta) d\Theta}{(\varphi(y))^{\frac{mp-\beta}{1+p}}} \leq \frac{m(1+p)}{mp-\beta} \left( \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}} - \varepsilon \right)^{-\frac{1}{p}}.$$

By (3.19) we know that

$$m \int_0^{\varphi(y)} \Theta^{m-1} \Upsilon^{-\frac{1}{p}}(\Theta) d\Theta = y. \quad (3.20)$$

Using this fact and raising the last inequality to the power  $-\frac{1+p}{mp-\beta}$  we have

$$\left( \frac{m(1+p)}{mp-\beta} \left( \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}} - \varepsilon \right)^{-\frac{1}{p}} \right)^{-\frac{1+p}{mp-\beta}} \leq y^{-\frac{1+p}{mp-\beta}} \varphi(y) \leq \left( \frac{m(1+p)}{mp-\beta} \left( \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}} + \varepsilon \right)^{-\frac{1}{p}} \right)^{-\frac{1+p}{mp-\beta}}.$$

Note that the limit as  $y \rightarrow +\infty$  may not exist for the inequality above, but  $\liminf$  and  $\limsup$  do exist as  $y \rightarrow +\infty$ . So we have

$$\begin{aligned} \left( \frac{m(1+p)}{mp-\beta} \left( \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}} + \varepsilon \right)^{-\frac{1}{p}} \right)^{-\frac{1+p}{mp-\beta}} &\leq \liminf_{y \rightarrow +\infty} y^{-\frac{1+p}{mp-\beta}} \varphi(y) \leq \limsup_{y \rightarrow +\infty} y^{-\frac{1+p}{mp-\beta}} \varphi(y) \leq \\ &\left( \frac{m(1+p)}{mp-\beta} \left( \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}} - \varepsilon \right)^{-\frac{1}{p}} \right)^{-\frac{1+p}{mp-\beta}}. \end{aligned}$$

Now, passing  $\varepsilon \rightarrow 0^+$ , we have

$$\lim_{y \rightarrow +\infty} y^{-\frac{1+p}{mp-\beta}} \varphi(y) = \left( \frac{m(1+p)}{mp-\beta} \left( \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}} \right)^{-\frac{1}{p}} \right)^{-\frac{1+p}{mp-\beta}}.$$

Simplifying we have that

$$\begin{aligned} \left( \frac{m(1+p)}{mp-\beta} \left( \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{p}{1+p}} \right)^{-\frac{1}{p}} \right)^{-\frac{1+p}{mp-\beta}} &= \left[ \frac{bm(1+p)}{p(m+\beta)} \right]^{\frac{1}{mp-\beta}} \left[ \frac{m(1+p)}{mp-\beta} \right]^{-\frac{1+p}{mp-\beta}} = \\ &\left[ \frac{b(mp-\beta)^{1+p}}{(m(1+p))^p p(m+\beta)} \right]^{\frac{1}{mp-\beta}} = C_*. \end{aligned}$$

So, in fact, we have that

$$\lim_{y \rightarrow +\infty} y^{-\frac{1+p}{mp-\beta}} \varphi(y) = C_*,$$

which proves formula 2.

Now we prove formula 4., the proof of formula 3. is similar. From Lemma 3.1.4 we know that

$$\Upsilon(\Theta) \sim k\Theta, \text{ as } \Theta \rightarrow 0^+ \text{ if } k > 0, p(m+\beta) > 1+p.$$

Hence, for any  $\varepsilon > 0$ , there exists  $\Theta_\varepsilon > 0$  such that

$$(k-\varepsilon)\Theta \leq \Upsilon(\Theta) \leq (k+\varepsilon)\Theta, \quad \Theta \leq \Theta_\varepsilon,$$

$$(k-\varepsilon)(\varphi(y)) \leq ((\varphi^m(y))')^p \leq (k+\varepsilon)(\varphi(y))^{\frac{p(m+\beta)}{1+p}}, \quad \varphi(y) \leq \Theta_\varepsilon,$$

$$(k-\varepsilon)^{1/p}(\varphi(y))^{\frac{1}{p}} \leq m\varphi^{m-1}\varphi'(y) \leq (k+\varepsilon)^{1/p}(\varphi(y))^{\frac{1}{p}}, \quad \varphi(y) \leq \Theta_\varepsilon,$$

$$(k-\varepsilon)^{1/p}dy \leq m(\varphi(y))^{m-1-\frac{1}{p}}d\varphi \leq (k+\varepsilon)^{1/p}dy, \quad \varphi(y) \leq \Theta_\varepsilon.$$

Integrating we get

$$(k - \varepsilon)^{1/p} y \leq \frac{mp}{mp-1} (\varphi(y))^{\frac{mp-1}{p}} \leq (k + \varepsilon)^{1/p} y, \quad \varphi(y) \leq \Theta_\varepsilon,$$

$$\begin{aligned} \left[ \frac{mp-1}{mp} (k - \varepsilon)^{1/p} \right]^{\frac{p}{mp-1}} &\leq y^{-\frac{p}{mp-1}} \varphi(y) \leq \\ &\leq \left[ \frac{mp-1}{mp} (k + \varepsilon)^{1/p} \right]^{\frac{p}{mp-1}}, \quad \varphi(y) \leq \Theta_\varepsilon. \end{aligned}$$

So we have

$$\begin{aligned} \left[ \frac{mp-1}{mp} (k - \varepsilon)^{1/p} \right]^{\frac{p}{mp-1}} &\leq \liminf_{y \rightarrow 0^+} y^{-\frac{p}{mp-1}} \varphi(y) \leq \\ &\leq \limsup_{y \rightarrow 0^+} y^{-\frac{p}{mp-1}} \varphi(y) \leq \left[ \frac{mp-1}{mp} (k + \varepsilon)^{1/p} \right]^{\frac{p}{mp-1}}. \end{aligned}$$

Passing  $\varepsilon \rightarrow 0^+$  we have

$$\lim_{y \rightarrow 0^+} y^{-\frac{p}{mp-1}} \varphi(y) = \left[ \frac{mp-1}{mp} \right]^{\frac{p}{mp-1}} k^{\frac{1}{mp-1}},$$

which proves formula 4.

Finally, we prove formula 6. The proof of formula 5. is similar. From Lemma 3.1.4 we know that

$$\Upsilon(\Theta) \sim \left( -\frac{k}{bm} \right)^{-p} \Theta^{p(m+\beta-1)}, \text{ as } \Theta \rightarrow 0^+ \text{ if } k < 0, p(m+\beta) > 1+p.$$

Hence, for any  $\varepsilon > 0$ , there exists  $\Theta_\varepsilon > 0$  such that

$$\left( \left( -\frac{k}{bm} \right)^{-p} - \varepsilon \right) \Theta^{p(m+\beta-1)} \leq \Upsilon(\Theta) \leq \left( \left( -\frac{k}{bm} \right)^{-p} + \varepsilon \right) \Theta^{p(m+\beta-1)}, \quad \Theta \leq \Theta_\varepsilon,$$

$$\left( \left( -\frac{k}{bm} \right)^{-p} - \varepsilon \right) (\varphi(y))^{p(m+\beta-1)} \leq ((\varphi^m(y))')^p \leq \left( \left( -\frac{k}{bm} \right)^{-p} + \varepsilon \right) (\varphi(y))^{p(m+\beta-1)}, \quad \varphi(y) \leq \Theta_\varepsilon,$$



$$\begin{aligned} \left( \left( -\frac{k}{bm} \right)^{-p} - \varepsilon \right)^{1/p} (\varphi(y))^{(m+\beta-1)} &\leq m\varphi^{m-1}\varphi'(y) \leq \left( \left( -\frac{k}{bm} \right)^{-p} + \varepsilon \right)^{1/p} (\varphi(y))^{(m+\beta-1)}, \\ \left( \left( -\frac{k}{bm} \right)^{-p} - \varepsilon \right)^{1/p} dy &\leq m(\varphi(y))^{-\beta} d\varphi \leq \left( \left( -\frac{k}{bm} \right)^{-p} + \varepsilon \right)^{1/p} dy, \quad \varphi(y) \leq \Theta_\varepsilon. \end{aligned}$$

Integrating we get

$$\left( \left( -\frac{k}{bm} \right)^{-p} - \varepsilon \right)^{1/p} y \leq \frac{m}{1-\beta} (\varphi(y))^{1-\beta} \leq \left( \left( -\frac{k}{bm} \right)^{-p} + \varepsilon \right)^{1/p} y, \quad \varphi(y) \leq \Theta_\varepsilon,$$

$$\begin{aligned} \left[ \frac{1-\beta}{m} \left( \left( -\frac{k}{bm} \right)^{-p} - \varepsilon \right)^{1/p} \right]^{\frac{1}{1-\beta}} &\leq y^{-\frac{1}{1-\beta}} \varphi(y) \leq \\ &\leq \left[ \frac{1-\beta}{m} \left( \left( -\frac{k}{bm} \right)^{-p} + \varepsilon \right)^{1/p} \right]^{\frac{1}{1-\beta}}, \quad \varphi(y) \leq \Theta_\varepsilon. \end{aligned}$$

So we have

$$\begin{aligned} \left[ \frac{1-\beta}{m} \left( \left( -\frac{k}{bm} \right)^{-p} - \varepsilon \right)^{1/p} \right]^{\frac{1}{1-\beta}} &\leq \liminf_{y \rightarrow 0^+} y^{-\frac{1}{1-\beta}} \varphi(y) \leq \\ &\leq \limsup_{y \rightarrow 0^+} y^{-\frac{1}{1-\beta}} \varphi(y) \leq \left[ \frac{1-\beta}{m} \left( \left( -\frac{k}{bm} \right)^{-p} + \varepsilon \right)^{1/p} \right]^{\frac{1}{1-\beta}}. \end{aligned}$$

Passing  $\varepsilon \rightarrow 0^+$  we have

$$\lim_{y \rightarrow 0^+} y^{-\frac{1}{1-\beta}} \varphi(y) = \left( (1-\beta) \left( -\frac{b}{k} \right) \right)^{\frac{1}{1-\beta}},$$

which proves formula 6.

Note that letting  $y \rightarrow \ell^-$  in (3.20), we obtain  $\ell = +\infty$ , if and only if

$$\int_0^{+\infty} \Upsilon^{-\frac{1}{p}}(z) dz = +\infty. \quad (3.21)$$

By Lemma 3.1.4, we know that the integral above (3.21) is true, then the solution to the problem (3.8) is global. It follows that there exists a finite traveling-wave solution of the

form:  $u(x, t) = \varphi(kt - x)$ , to equation (1.7).

□

## Chapter 4

# Evolution of Interfaces for the Nonlinear Double Degenerate Parabolic Equation with Fast Diffusion

We present a full qualitative analysis of the asymptotic behavior of the interfaces (when they exist) and local asymptotic behavior of the solution near them for the problem (1.7), (1.9) in the fast diffusion case ( $0 < mp < 1$ ) in this chapter. In the cases where there is an infinite speed of propagation ( $\eta(t) = +\infty$ ), we classify the asymptotic behavior of the solution at infinity. The results of Chapter 4 have recently been submitted for publication, [24].

### 4.1 Main Results

We assume that  $u$  is a unique weak solution of the CP (1.7), (1.9), throughout this section. The main results are classified according to regions I-V, respectively, in Figure 4.1,

the  $(\alpha, \beta)$  parameter space diagram, below. We refer to Appendix 6.2 for the explicit values of relevant constants that appear throughout this section.




Figure 4.1:  $(\alpha, \beta)$  parameter space diagram for the interface development for the CP (1.7), (1.9) with  $0 < mp < 1$ , which is presented in [24].

- **Region I:**  $b > 0, 0 < \beta < mp$  and  $0 < \alpha < (1+p)/(mp-\beta)$ .

The interface initially expands and there exists a number  $\delta > 0$  such that

$$z_1 t^{\frac{mp-\beta}{(1-\beta)(1+p)}} \leq \eta(t) \leq z_2 t^{\frac{mp-\beta}{(1-\beta)(1+p)}}, \quad 0 \leq t \leq \delta, \quad (4.1)$$

Further, for any  $\sigma \in \mathbb{R}$ , there is a number  $f(\sigma) > 0$  (depending on  $C, m$ , and  $p$ ) such that

$$u(\chi_\sigma(t), t) \sim f(\sigma) t^{\frac{\alpha}{1+p-\alpha(mp-1)}}, \quad \text{as } t \rightarrow 0^+, \quad (4.2)$$

where  $\chi_\sigma(t) = \sigma t^{\frac{1}{1+p-\alpha(mp-1)}}$ .

- **Region II:**  $b > 0, 0 < \beta < mp, \alpha = (1 + p)/(mp - \beta)$ , and

$$C_* = \left[ \frac{b(mp - \beta)^{1+p}}{(m(1 + p))^p p(m + \beta)} \right]^{\frac{1}{mp - \beta}}.$$

Then the interface shrinks or expands accordingly as  $C < C_*$  or  $C > C_*$  and we have that

$$\eta(t) \sim z_* t^{\frac{mp - \beta}{(1+p)(1-\beta)}}, \text{ as } t \rightarrow 0^+, \quad (4.3)$$

where  $z_* \leq 0$  if  $C \leq C_*$ , and for any  $\sigma < z_*$ , there is a number  $f_1(\sigma) > 0$  such that

$$u(z_\sigma(t), t) \sim t^{1/(1-\beta)} f_1(\sigma), \text{ as } t \rightarrow 0^+, \quad (4.4)$$

where  $z_\sigma(t) = \sigma t^{\frac{mp - \beta}{(1-\beta)(1+p)}}$ .

- **Region III:**  $b > 0, 0 < \beta < mp$ , and  $\alpha > (1 + p)/(mp - \beta)$ .

Then the interface initially shrinks and we have that

$$\eta(t) \sim -\tau_* t^{\frac{1}{\alpha(1-\beta)}}, \text{ as } t \rightarrow 0^+, \quad (4.5)$$

where  $\tau_* = C^{-1/\alpha} (b(1 - \beta))^{\frac{1}{\alpha(1-\beta)}}$  and, for any  $\tau > \tau_*$ , we have

$$u(\eta_\tau(t), t) \sim \left[ C^{1-\beta} \tau^{\alpha(1-\beta)} - b(1 - \beta) \right]^{\frac{1}{1-\beta}} t^{\frac{1}{1-\beta}}, \text{ as } t \rightarrow 0^+, \quad (4.6)$$

where  $\eta_\tau(t) = -\tau t^{\frac{1}{\alpha(1-\beta)}}$ .

- **Region IV:**  $b > 0, 0 < \beta = mp < 1$ , and  $\alpha > 0$ .

In this case the solution travels with an infinite speed of propagation: there is no interface. For arbitrary  $\varepsilon > 0$ , there exists a number  $\delta = \delta(\varepsilon) > 0$  such that

$$t^{\frac{1}{1-mp}} \vartheta(x) \leq u(x, t) \leq (t + \varepsilon)^{\frac{1}{1-mp}} \vartheta(x), \quad x > 0, \quad 0 \leq t \leq \delta, \quad (4.7)$$

where  $\vartheta = \vartheta(x) > 0$  is a solution of the stationary problem

$$\begin{cases} ((\vartheta^m)'|^{p-1}(\vartheta^m)')' - \frac{1}{1-mp}\vartheta - b\vartheta^{mp} = 0, & x > 0, \\ \vartheta(0) = 1, \vartheta(+\infty) = 0. \end{cases} \quad (4.8)$$

Further, we have the estimation

$$\ln u(x, t) \sim -\frac{1}{m} \left(\frac{b}{p}\right)^{1/(1+p)} x, \quad \text{as } x \rightarrow +\infty, \quad 0 \leq t \leq \delta. \quad (4.9)$$

- **Region V:** This region is divided into cases V(a), V(b) and V(b).

– **V(a):** Either  $b > 0, \beta > mp$  or  $b < 0, \beta \geq 1$  or  $b = 0$ , and

$$\mathcal{D} = \left[ \frac{(m(1+p))^p (m+1)}{(1-mp)^p} \right]^{\frac{1}{1-mp}}.$$

In this case the solution travels with an infinite speed of propagation: there is no interface; and (4.2) holds. If  $b > 0, \beta \geq \frac{p(1-m)+2}{1+p}$  or  $b < 0, \beta \geq 1$  or  $b = 0$ , then there exists a number  $\delta > 0$  such that

$$u(x, t) \sim \mathcal{D} t^{\frac{1}{1-mp}} x^{\frac{1+p}{1-mp}}, \quad \text{as } x \rightarrow +\infty, \quad t \in (0, \delta], \quad (4.10)$$

– **V(b)**:  $b > 0$  and  $1 \leq \beta < \frac{p(1-m)+2}{1+p}$ .

Then,

$$\lim_{t \rightarrow 0^+} \lim_{x \rightarrow +\infty} \frac{u(x, t)}{t^{\frac{1}{1-mp}} x^{\frac{1+p}{mp-1}}} = \mathcal{D}. \quad (4.11)$$

– **V(c)**:  $b > 0$  and  $0 < mp < \beta < 1$ .

Then

$$u(x, t) \sim C_* x^{\frac{1+p}{mp-\beta}}, \text{ as } x \rightarrow +\infty, \quad (4.12)$$

holds for any  $t \in (0, \delta]$  for some  $\delta$ .

## 4.2 Additional Details of the Results

In this section we outline some additional, essential details of Results I-V. We refer to Appendix 6.2 for the explicit values of relevant constants that appear throughout this section.

- *Region I.* The solution  $u$  satisfies the following estimation

$$C_1 t^{\frac{1}{1-\beta}} (z_1 - z)_+^{\frac{1+p}{mp-\beta}} \leq u(x, t) \leq C_* t^{\frac{1}{1-\beta}} (z_2 - z)_+^{\frac{1+p}{mp-\beta}}, \quad 0 < t \leq \delta, \quad (4.13)$$

where  $z = xt^{\frac{\beta-mp}{(1-\beta)(1+p)}}$ . The left-hand side of (4.13) is valid for  $0 \leq x < +\infty$  and the right-hand side is valid in the curved region:  $x \geq \tau_0 t^{(mp-\beta)/(1-\beta)(1+p)}$ .  $C_1, z_1, z_2,$

and  $\tau_0$ , are positive constants depending on  $m, p, \beta$ , and  $b$ . Moreover, the function  $f$  is such that

$$f(\sigma) = C^{\frac{1+p}{1+p-\alpha(mp-1)}} f_0(C^{\frac{mp-1}{1+p-\alpha(mp-1)}} \sigma), \quad f_0(\sigma) = w(\sigma, 1), \quad \sigma \in \mathbb{R}, \quad (4.14)$$

where  $w$  is a minimal solution of the CP (1.7), (1.10) with  $C = 1$  and  $b = 0$ . If  $u_0$  is given by (1.10), then the right-hand sides of (4.13) and (4.1) are valid for all  $t > 0$ .

- *Region II.* Assume  $u$  solves the CP (1.7), (1.10). If  $C = C_*$  in  $u_0$ , then  $u = u_0$  is the stationary solution to the CP. If  $C \neq C_*$ , then the minimal solution of the CP is given by

$$u(x, t) = t^{\frac{1}{1-\beta}} f_1(z), \quad z = xt^{\frac{\beta-mp}{(1-\beta)(1+p)}}, \quad (4.15)$$

and

$$\eta(t) = z_* t^{\frac{mp-\beta}{(1-\beta)(1+p)}}, \quad t \geq 0, \quad (4.16)$$

If  $C > C_*$ , the interface initially expands and we have the following estimation

$$C' \left( z' t^{\frac{mp-\beta}{(1-\beta)(1+p)}} - x \right)_+^{\frac{1+p}{mp-\beta}} \leq u \leq C'' \left( z'' t^{\frac{mp-\beta}{(1-\beta)(1+p)}} - x \right)_+^{\frac{1+p}{mp-\beta}}, \quad (4.17)$$

$$z' \leq z_* \leq z'', \quad 0 \leq x < +\infty, \quad t > 0, \quad (4.18)$$

where  $C' = C_2, C'' = C_*, z' = z_3$ , and  $z'' = z_4$ . While, if  $0 < C < C_*$ , then the interface shrinks and there exists a number  $\tau_1 > 0$  such that for all  $\tau \leq \tau_1$  there exists a number  $\varrho > 0$  such that

$$u\left(\tau t^{\frac{mp-\beta}{(1+p)(1-\beta)}}, t\right) = \varrho t^{\frac{1}{1-\beta}}, \quad t \geq 0, \quad (4.19)$$



and  $u$  and  $z_*$  satisfy the estimates (4.17) and (4.18) with  $C' = C_*$ ,  $C'' = C_3$ ,  $z' = -z_5$ , and  $z'' = -z_6$ .

- *Region IV.* The explicit solution of the problem (4.8) is given by the following formula

$$\vartheta(x) = \mathcal{J}^{-1}(x), \quad 0 \leq x < +\infty, \quad (4.20)$$

where  $\mathcal{J}^{-1}(\cdot)$  is the inverse function of

$$\mathcal{J}(z) = \int_z^1 ms^{-1} \left[ \frac{b}{p} + \frac{m(1+p)}{p(1-mp)(1+m)} s^{1-mp} \right]^{-\frac{1}{1+p}} ds, \quad 0 < z \leq 1. \quad (4.21)$$

The function  $\vartheta(x)$  satisfies the asymptotic formula

$$\ln \vartheta(x) \sim -\frac{1}{m} \left( \frac{b}{p} \right)^{1/(1+p)} x, \quad \text{as } x \rightarrow +\infty, \quad (4.22)$$

and the global estimation

$$0 < \vartheta(x) \leq \exp \left( -\frac{1}{m} \left( \frac{b}{p} \right)^{\frac{1}{1+p}} x \right), \quad x > 0. \quad (4.23)$$

From (4.23), it follows that

$$\frac{\vartheta(x)}{e^{-\gamma x}} \rightarrow +\infty, \quad \text{as } x \rightarrow +\infty, \quad \text{if } \gamma > \frac{1}{m} \left( \frac{b}{p} \right)^{\frac{1}{1+p}}. \quad (4.24)$$

From (4.7) and (4.24), it follows that

$$\lim_{t \rightarrow 0^+} \lim_{x \rightarrow +\infty} u(x, t) \exp \left( -\frac{1}{m} \left( \frac{b}{p} \right)^{\frac{1}{1+p}} x \right) = 0, \quad (4.25)$$

and respectively

$$\frac{u(x, t)}{e^{-\gamma x}} \rightarrow +\infty, \text{ as } x \rightarrow +\infty, 0 \leq t \leq \delta(\varepsilon), \text{ if } \gamma > \frac{1}{m} \left( \frac{b}{p} \right)^{\frac{1}{1+p}}. \quad (4.26)$$

- *Region V.* If  $\beta \geq 1$ , then for arbitrary  $\varepsilon > 0$ , there exists  $\delta = \delta(\varepsilon) > 0$  such that we have the following estimation

$$C_5 t^{\frac{\alpha}{1+p-\alpha(mp-1)}} (\chi_1 + \chi)^{\frac{1+p}{mp-1}} \leq u(x, t) \leq C_6 t^{\frac{\alpha}{1+p-\alpha(mp-1)}} (\chi_2 + \chi)^{\frac{1+p}{mp-1}}, \quad (4.27)$$

where  $\chi = xt^{\frac{-1}{1+p-\alpha(mp-1)}}$ , for all  $x \in [0, \infty)$  and  $0 \leq t \leq \delta(\varepsilon)$ .  $C_5, C_6, \chi_1$ , and  $\chi_2$ , are positive constants depending on  $m, p, \beta, b$ , and  $\varepsilon$ .

If  $b > 0$  and  $\beta \geq 1$ , we have the upper estimation

$$u(x, t) \leq \mathcal{D} t^{\frac{1}{1-mp}} x^{\frac{1+p}{mp-1}}, 0 < x < +\infty, 0 < t < +\infty. \quad (4.28)$$

While if  $b < 0$  and  $\beta \geq 1$ , then for arbitrary  $\varepsilon > 0$ , there exists  $\delta = \delta(\varepsilon) > 0$  such that the following estimation is satisfied

$$u(x, t) \leq \mathcal{D}(1 - \varepsilon)^{\frac{1}{mp-1}} t^{\frac{1}{1-mp}} x^{\frac{1+p}{mp-1}}, \text{ for } \kappa t^{\frac{1}{1+p+\alpha(1-mp)}} < x < +\infty, 0 < t \leq \delta, \quad (4.29)$$

where

$$\kappa = \left[ \frac{A_0 + \varepsilon}{\mathcal{D}(1 - \varepsilon)^{\frac{1}{mp-1}}} \right]^{\frac{mp-1}{1+p}}. \quad (4.30)$$

If  $b > 0$  and  $mp < \beta < 1$ , then there exists  $\delta > 0$  such that

$$t^{\frac{1}{1-\beta}} C_* (1 - \varepsilon)(z_8 + z)^{\frac{1+p}{mp-\beta}} \leq u(x, t) \leq C_* x^{\frac{1+p}{mp-\beta}}, 0 < x < +\infty, 0 < t \leq \delta, \quad (4.31)$$

where  $z = xt^{\frac{\beta-mp}{(1-\beta)(1+p)}}$ ,  $\varepsilon > 0$  is an arbitrary sufficiently small number, and  $z_8$  is a positive constant depending on  $m, p, \beta, b$ , and  $\varepsilon$ .

If  $b = 0$  and  $\alpha > 0$ , then the minimal solution to the CP (1.7), (1.10) has the self-similar form

$$u(x, t) = t^{\frac{\alpha}{1+p+\alpha(1-mp)}} f(\chi), \chi = xt^{\frac{-1}{1+p+\alpha(1-mp)}}. \quad (4.32)$$

where  $f$  satisfies (4.14). Further, following global estimation is valid

$$\mathcal{D}t^{\frac{\alpha}{1+p+\alpha(1-mp)}} (\chi_3 + \chi)^{\frac{1+p}{mp-1}} \leq u(x, t) \leq C_7 t^{\frac{\alpha}{1+p+\alpha(1-mp)}} (\chi_4 + \chi)^{\frac{1+p}{mp-1}}, \quad (4.33)$$

$$0 \leq x < +\infty, 0 < t < +\infty,$$

where  $C_7, \chi_3$ , and  $\chi_4$ , are positive constants depending on  $m$  and  $p$ .

Explicit solution (4.29) provides sharper upper bound than (4.33) as  $x \rightarrow +\infty$ .

From (4.33) and (4.29), asymptotic result (4.10) easily follows. In a similar way asymptotic result (4.10) follows in the local case, (1.9).

## 4.3 Asymptotic Properties of Solutions With $0 < mp < 1$

In this section we establish a series of lemmas that describe preliminary estimations for the CP. The proof of these results is based on nonlinear scaling.

### 4.3.0.1 Diffusion Weakly Dominates Reaction

**Lemma 4.3.1.** *If  $b = 0$  and  $\alpha > 0$ , then the global formula (4.32) holds,*

$$u(x, t) = t^{\alpha/(1+p-\alpha(mp-1))} f(\chi), \chi = xt^{-\frac{1}{1+p-\alpha(mp-1)}},$$

where  $u$  is a minimal solution of the CP (1.7), (1.10) and the self-similarity function  $f$  satisfies (4.14),

$$\begin{cases} (|(f^m(\chi))'|^{p-1}(f^m(\chi))')' + (1+p-\alpha(mp-1))^{-1}(\chi f'(\chi) - \alpha f(\chi)) = 0, \\ f(\chi) \sim C(-\chi)^\alpha, \text{ as } \chi \downarrow -\infty, \text{ and } f(\chi) \rightarrow 0, \text{ as } \chi \uparrow +\infty. \end{cases}$$

If  $u_0$  satisfies (1.9), and  $u$  is the unique weak solution to CP (1.7), (1.8), then  $u$  satisfies (4.2).

The proof coincides with that given for Lemma 2.3.4.

**Lemma 4.3.2.** *Let  $u$  be a weak solution to the CP (1.7), (1.8), with  $u_0$  satisfying the condition (1.9). Let one of the following cases be valid*

$$\begin{cases} b > 0, 0 < \beta < mp, 0 < \alpha < (1+p)/(mp-\beta) & \text{Case 1,} \\ b > 0, \beta \geq mp, \alpha > 0 & \text{Case 2,} \\ b < 0, \beta \geq 1, \alpha > 0 & \text{Case 3.} \end{cases}$$

Then, for any  $\sigma \in \mathbb{R}$ ,  $u$  satisfies (4.2) with the same function  $f$  as in Lemma 4.3.1.

The proof coincides with the proof of Lemma 2.3.5. A minor change is required in order to prove Case 3. On the lines  $x = -x_\varepsilon$  and  $x = -k^{1/\alpha}x_\varepsilon$ , due to the infinite speed of propagation property, we choose

$$u_{\pm\varepsilon}(-x_\varepsilon, t) = u(-x_\varepsilon, t), 0 \leq t \leq \delta,$$

and

$$u_k^{\pm\varepsilon}(-k^{1/\alpha}x_\varepsilon, t) = u(-x_\varepsilon, k^{(\alpha(mp-1)-(1+p))/\alpha}, t), 0 \leq t \leq k^{(1+p-\alpha(mp-1))/\alpha}\delta,$$

rather than a zero boundary condition. The rest of the proof of Case 3 coincides with the proof of Lemma 2.3.5 when  $b < 0$ .

#### 4.3.0.2 Diffusion and Reaction are in Balance

**Lemma 4.3.3.** *If  $b > 0$ ,  $0 < \beta < mp < 1$ , and  $\alpha = (1 + p)/(mp - \beta)$ , then the minimal solution  $u$  to the CP (1.7), (1.9) has the self-similar form (4.15), where the self-similarity function  $f_1$  satisfies*

$$\begin{cases} \mathcal{L}^0 f_1 \equiv (|(f_1^m)'|^{p-1}(f_1^m)')' + \frac{mp-\beta}{(1+p)(1-\beta)} z f_1' - \frac{1}{1-\beta} f_1 - b f_1^\beta = 0, & z \in \mathbb{R}, \\ f_1(z) \sim C(-z)^{(1+p)/(mp-\beta)}, & \text{as } z \downarrow -\infty, \text{ and } f_1(z) \rightarrow 0, \text{ as } z \uparrow +\infty. \end{cases} \quad (4.34)$$

There exists  $\tau_1, \varrho > 0$  such that for any  $\tau \in (-\infty, -\tau_1)$  we have

$$u\left(\tau t^{\frac{mp-\beta}{(1+p)(1-\beta)}}, t\right) = \varrho t^{\frac{1}{1-\beta}}, \quad t \geq 0. \quad (4.35)$$

If  $0 < C < C_*$ , then we have

$$0 < \varrho < C_* (-\tau)^{\frac{1+p}{mp-\beta}}, \quad (4.36)$$

while if  $C > C_*$ , then  $f_1(0) = A_1(m, p, \beta, C, b) = A_1 > 0$ .

*Proof of Lemma 4.3.3.* We define

$$u_k(x, t) = ku(k^{\frac{\beta-mp}{1+p}} x, k^{\beta-1} t), \quad k > 0, \quad (4.37)$$

(see Lemma 6 of [20]). The function (4.37) satisfies the CP (1.7), (1.10). We consider  $u$  to be a unique minimal solution of CP (1.7), (1.10) such that

$$u(x, t) \leq ku(k^{\frac{\beta-mp}{1+p}} x, k^{\beta-1} t), \quad k > 0. \quad (4.38)$$

By changing the variables in (4.38) as

$$y = k^{\frac{\beta-mp}{1+p}} x, \ell = k^{\beta-1} t, \quad (4.39)$$

we derive (4.38) with opposite inequality and with  $k$  replaced with  $k^{-1}$ . Since  $k > 0$  is arbitrary, (4.38) follows with "=". Taking  $k = t^{1/(1-\beta)}$ , (4.37) implies (4.15) with  $f_1(z) = u(z, 1)$ , where  $f_1$  is a solution to problem (4.34). This proves the first part of the lemma. The second part of the lemma is proved in the same way as the second part of Lemma 3.3 of [6].  $\square$

**Lemma 4.3.4.** *Let  $b > 0$ ,  $0 < \beta < mp < 1$ , and  $\alpha = (1+p)/(mp-\beta)$ , and let  $u$  be the minimal solution to the CP (1.7), (1.9). Then  $u$  satisfies*

$$u\left(\tau t^{\frac{mp-\beta}{(1+p)(1-\beta)}}, t\right) \sim \varrho t^{\frac{1}{1-\beta}}, \text{ as } t \rightarrow 0^+, \quad (4.40)$$

where  $\tau_1, \varrho > 0$  are the same as in Lemma 4.3.3. Furthermore, if  $0 < C < C_*$ , then

$$0 < \varrho < C_* (-\tau)^{\frac{1+p}{mp-\beta}}. \quad (4.41)$$

If  $C > C_*$ , then

$$u(0, t) \sim A_1 t^{\frac{1}{1-\beta}}, \text{ as } t \rightarrow 0^+; f_1(0) = A_1 > 0. \quad (4.42)$$

The proof of Lemma 4.3.4 follows as a localization of the proof of Lemma 4.3.3, exactly as local results were proven for Lemma 2.3.5.

### 4.3.0.3 Reaction Dominates Diffusion

**Lemma 4.3.5.** *If  $b > 0$ ,  $0 < \beta < mp < 1$ , and  $\alpha > (1+p)/(mp-\beta)$ , then the unique weak solution  $u$  to the CP (1.7), (1.9) satisfies (4.6).*

The proof of Lemma 4.3.5 coincides with the proof of Lemma 2.3.8.

## 4.4 Proofs of the Main Results

From Lemma 4.3.2, the asymptotic formulas (4.2) and (4.14) follow. For any  $\varepsilon > 0$ , from (4.2), we have

$$(A_0 - \varepsilon)t^{\alpha/(1+p-\alpha(mp-1))} \leq u(0, t) \leq (A_0 + \varepsilon)t^{\alpha/(1+p-\alpha(mp-1))}, \quad 0 \leq t \leq \delta_1(\varepsilon), \quad (4.43)$$

for some  $\delta_1 = \delta_1(\varepsilon) > 0$  and where  $A_0 = f(0) > 0$ .

The proof of results for Region I follows exactly as the proof of result (1) from [6] for  $b \neq 0$ , by choosing

$$r(x, t) = t^{\frac{1}{1-\beta}} f_1(z), \quad z = xt^{\frac{\beta-mp}{(1-\beta)(1+p)}}, \quad (4.44)$$

$$f_1 = C_0(z_0 - z)_+^{\frac{1+p}{mp-\beta}}, \quad 0 < z < +\infty \quad (4.45)$$

with  $C_0, z_0 > 0$  to be determined. To prove the left-hand sides of estimates (4.13) and (4.1), we choose  $C_0 = C_1$  and  $z_0 = z_1$  (see the Appendix 6.2) and apply comparison theorem. We prove the right-hand sides of the estimations (4.13) and (4.1) by choosing  $C_0 = C_*$ ,  $z_0 = z_2$  and  $\tau_0$  and applying comparison theorem in the curved region  $G_{\tau_0, \delta}$ , where

$$G_{\tau, \delta} = \{(x, t) : z_\tau(t) = \tau t^{\frac{mp-\beta}{(1+p)(1-\beta)}} < x < +\infty, 0 < t \leq \delta\}.$$

*Region II.* Assume that  $u_0$  is defined as (1.10). The self-similar solution (4.15) follows from Lemma 4.3.3. The proof of estimation (4.17) when  $C > C_*$  (also when  $u_0$  is given through (1.9)) coincides with the proof given in [20]. Let  $0 < C < C_*$ . The formula (4.19) follows from Lemma 4.3.3. The proof of the right-hand side of (4.17) (also when  $u_0$  is given through (1.9)) coincides with the proof given in [20] and the proof of the left-hand side of (4.17) follows in the same way as the proof of the analogous estimate from result (3) of [6].  $\square$

*Region III.* The asymptotic estimate (4.6) follows from Lemma 4.3.4. The proof of the asymptotic estimate (4.5) coincides with the proof of Theorem 2.1.3.  $\square$

*Region IV.* The asymptotic estimation (4.2) is proved in Lemma 4.3.2. From (4.2), (4.43) follows. The proof of estimate (4.7) follows in the same way as the analogous estimate in result (4) from [6].

Intergration of (4.8) implies (4.20). By rescaling  $x$  with  $\varepsilon^{-1}x$ ,  $\varepsilon > 0$  from (4.20) we have

$$\frac{x}{\varepsilon} = \int_{\vartheta(\frac{x}{\varepsilon})}^1 \frac{m}{s} \left[ \frac{b}{p} + \frac{m(1+p)}{p(1-mp)(1+m)} s^{1-mp} \right]^{-\frac{1}{1+p}} ds, \quad s > 0.$$

Letting  $r = -\varepsilon \ln s$  implies

$$x = \mathcal{J}[\Lambda_\varepsilon(x)], \quad (4.46)$$

where

$$\mathcal{J}(s) = \int_0^s m \left[ \frac{b}{p} + \frac{m(1+p)}{p(1-mp)(1+m)} e^{r(mp-1)/\varepsilon} \right]^{-\frac{1}{1+p}} dr,$$

and

$$\Lambda_\varepsilon(x) = -\varepsilon \ln \vartheta\left(\frac{x}{\varepsilon}\right).$$

From (4.46) it follows that

$$\Lambda_\varepsilon(x) = \mathcal{J}^{-1}(x), \quad (4.47)$$



where  $\mathcal{J}^{-1}$  denotes the inverse function of  $\mathcal{J}$ .

Since  $0 < mp < 1$  it follows that

$$\lim_{\varepsilon \rightarrow 0^+} \mathcal{J}(y) = m(b/p)^{-\frac{1}{1+p}} y, \quad \lim_{\varepsilon \rightarrow 0^+} \mathcal{J}^{-1}(y) = m^{-1}(b/p)^{\frac{1}{1+p}} y, \quad (4.48)$$

for  $y \geq 0$ , uniformly, on bounded subsets. From (4.47) and (4.48) it follows that

$$-\lim_{\varepsilon \rightarrow 0^+} \varepsilon \ln \vartheta\left(\frac{x}{\varepsilon}\right) = m^{-1}(b/p)^{\frac{1}{1+p}} x. \quad (4.49)$$

The asymptotic formula (4.22) follows by choosing  $y = x/\varepsilon$ . Inequality (4.23), as well as estimations (4.24),(4.25),(4.26) follow from (4.20) and (4.21).  $\square$

*Region V.* Let either  $b > 0, \beta > mp$  or  $b < 0, \beta \geq 1$ . The proof of this case follows in the same way as the analogous case of result (5) of [6] Formula (4.2) is a consequence of Lemma 4.3.2. Choose  $\varepsilon > 0$ , sufficiently small. From (4.2), (4.43) holds for some  $\delta_1 = \delta_1(\varepsilon) > 0$ . Let  $\beta \geq 1$ . Consider

$$r(x, t) = t^{\frac{\alpha}{1+p+\alpha(1-mp)}} f(\chi), \chi = xt^{\frac{-1}{1+p+\alpha(1-mp)}}. \quad (4.50)$$

We have

$$Lr = t^{\frac{\alpha mp - 1 - p}{1+p+\alpha(1-mp)}} L_1 f, \quad (4.51)$$

where

$$L_1 f = \frac{\alpha}{1+p+\alpha(1-mp)} f - \frac{1}{1+p+\alpha(1-mp)} \chi f' - \left( |(f^m)'|^{p-1} (f^m)' \right)' + bt^{\frac{1+p-\alpha(mp-\beta)}{1+p+\alpha(1-mp)}} f^\beta. \quad (4.52)$$

As a function  $f$  we select

$$f(\chi) = C_0(\chi_0 + \chi)^{\frac{1+p}{mp-1}}, \chi \geq 0, \quad (4.53)$$

where  $C_0$  and  $\chi_0$  are positive constants.

From (4.43) and Lemma 1 of [20], the right-hand side of (4.27) follows with  $\delta = \delta_2$ , such that

$$\delta_2 = \delta_1, \text{ if } b > 0$$

and

$$\delta_2 = \min(\delta_1, \delta_3), \text{ if } b < 0,$$

with

$$\delta_3 = \left[ \frac{\alpha \varepsilon (A_0 + \varepsilon)^{1-\beta}}{(1 + \varepsilon)(-b(1 + p + \alpha(1 - mp)))} \right]^{\frac{1+p+\alpha(1-mp)}{1+p+\alpha(\beta-mp)}}.$$

To prove a lower bound in this case we take  $C_0 = C_5$  and  $\chi_0 = \chi_1$ .

The proof of the left-hand side of (4.27) if either  $b > 0, \beta < \frac{p(1-m)+2}{1+p}$  or  $b < 0, \beta \geq 1$  or  $b > 0, \beta \geq \frac{p(1-m)+2}{1+p}$ , follows in the same way as the analogous estimate in result (5) of [6].

If  $b > 0$  and  $\beta \geq 1$ , then the proof of estimates (4.10), (4.11), and (4.29) follow as the analogous proof from result (5) of [6]. While if  $b > 0$  and  $0 < mp < \beta < 1$  the left-hand side of (4.31) follows as left-hand side of (4.13) was previously proved by choosing  $f_1(z) = C_*(1 - \varepsilon)(z_8 + z)_+^{\frac{1+p}{mp-\beta}}$  in (4.44). The proof of (4.12) follows in the same way as the proof of the analogous result from result (5) of [6].

Now, let  $b = 0$ . Let  $u_0$  be given by (1.10). Formula (4.32) is a consequence of Lemma 4.3.1. In order to prove inequality (4.33), choose the function  $g$  as in (4.50); satisfying (4.51) with  $b = 0$ . Choose the function  $f$  as (4.53). The proof of estimate (4.33) follows

in the same way as the proof of the analogous result from result (5) of [6]. To prove an upper estimate we choose  $C_0 = C_7$  and  $\chi_0 = \chi_4$  and to prove a lower estimation we choose  $C_0 = \mathcal{D}$  and  $\chi_0 = \chi_3$ . □

# Chapter 5

## Conclusions

A full qualitative analysis of the short-time behavior of the interface function and local solution near the interface function or at infinity (when the interface does not exist) for the Cauchy problem for the nonlinear double degenerate parabolic equation of turbulent filtration with absorption

$$u_t = (|(u^m)_x|^{p-1}(u^m)_x)_x - bu^\beta, \quad x \in \mathbb{R}, \quad 0 < t < T, \quad \beta > 0,$$

with

$$u(x, 0) = u_0(x) \sim C(-x)_+^\alpha, \quad \text{as } x \rightarrow 0^-, \quad \text{for some } C > 0, \quad \alpha > 0,$$

is successfully pursued in this dissertation in the case of both slow ( $mp > 1$ ) and fast ( $0 < mp < 1$ ) diffusion with either  $b \geq 0$  with  $0 < \beta < 1$  or  $b < 0$  with  $\beta \geq 1$ .

The following is a summary of the main results.

Slow Diffusion Case:  $mp > 1$ .

- If  $b > 0$ ,  $0 < \beta < 1$ , and  $0 < \alpha < (1 + p)/(mp - \beta)$ , then the diffusion term dominates over the reaction term and the interface expands forward, with the asymptotic formula for the interface function given by the following expression

$$\eta(t) \sim \psi t^{(mp-\beta)/(1-\beta)(1+p)}, \text{ as } t \rightarrow 0^+,$$

where  $\psi = \psi(C, m, p, \alpha) > 0$ .

- If  $b > 0$ ,  $0 < \beta < 1$ , and  $\alpha = (1 + p)/(mp - \beta)$ , then the diffusion and reaction terms are in balance, borderline case, and there is a critical constant  $C_*$ , such that the interface expands forward if  $C > C_*$  or shrinks backward if  $C < C_*$ , with the asymptotic formula for the interface function given by the following expression

$$\eta(t) \sim \zeta_* t^{(mp-\beta)/(1-\beta)(1+p)}, \text{ as } t \rightarrow 0^+,$$

where  $\zeta_* = \zeta_*(C, m, p) \leq 0$  if  $C \leq C_*$ .

- If  $b > 0$ ,  $0 < \beta < 1$ , and  $\alpha > (1 + p)/(mp - \beta)$ , then the reaction term *strongly* dominates over the diffusion term and the interface shrinks backwards, with the asymptotic formula for the interface function given by the following expression

$$\eta(t) \sim -\tau_* t^{1/\alpha(1-\beta)}, \text{ as } t \rightarrow 0^+,$$

where  $\tau_* = \tau_*(C, m, p, \alpha, \beta) > 0$ .

- If  $b \in \mathbb{R}$ ,  $\beta \geq 1$ , and  $\alpha \geq (1 + p)/(mp - 1)$ , then interface initially remains stationary.

Fast Diffusion Case:  $0 < mp < 1$ .

- If  $b > 0$ ,  $0 < \beta < mp$ , and  $0 < \alpha < (1 + p)/(mp - \beta)$ , then the diffusion term dominates over the reaction term and the interface expands forward, with the asymptotic formula for the interface function given by the following expression

$$\eta(t) \sim \psi t^{(mp-\beta)/(1-\beta)(1+p)}, \text{ as } t \rightarrow 0^+,$$

where  $\psi = \psi(C, m, p, \alpha) > 0$ .

- If  $b > 0$ ,  $0 < \beta < mp$ , and  $\alpha = (1 + p)/(mp - \beta)$ , then the diffusion and reaction terms are in balance, borderline case, and there is a critical constant  $C_*$ , such that the interface expands forward if  $C > C_*$  or shrinks backward if  $C < C_*$ , with the asymptotic formula for the interface function given by the following expression

$$\eta(t) \sim z_* t^{(mp-\beta)/(1-\beta)(1+p)}, \text{ as } t \rightarrow 0^+,$$

where  $z_* = z_*(C, m, p) \leq 0$  if  $C \leq C_*$ .

- If  $b > 0$ ,  $0 < \beta < mp$ , with  $\alpha > (1 + p)/(mp - \beta)$ , then the reaction term *strongly* dominates over the diffusion term and the interface shrinks backwards, with the asymptotic formula for the interface function given by the following expression

$$\eta(t) \sim -\tau_* t^{1/\alpha(1-\beta)}, \text{ as } t \rightarrow 0^+,$$

where  $\tau_* = \tau_*(C, m, p, \alpha, \beta) > 0$ .

- If either  $b > 0$ ,  $0 < \beta = mp < 1$ , and  $\alpha > 0$ , or  $b > 0$ ,  $\beta > mp > 0$  or  $b < 0$ ,  $\beta \geq 1$ , then the diffusion term *strongly* dominates over the reaction term and there is no interface ( $\eta(t) = +\infty$ ): the solution travels with an infinite speed of propagation.

If  $b > 0$ ,  $0 < \beta = mp < 1$  and  $\alpha > 0$  the solution decays as an exponential function at infinity; while if either  $b > 0, \beta > mp > 0$  or  $b < 0, \beta \geq 1$ , then the solution decays as a power-type function at infinity.

# Bibliography

- [1] U. G. Abdulla. Local structure of solutions of the Dirichlet problem for  $n$ -dimensional reaction-diffusion equations in bounded domains. *Advances in Differential Equations*, 4(2):197–224, 1999.
- [2] U. G. Abdulla. Reaction-diffusion in a closed domain formed by irregular curves. *Journal of Mathematical Analysis and Applications*, 246:480–492, 2000.
- [3] U. G. Abdulla. Reaction-diffusion in irregular domains. *Journal of Differential Equations*, 164(2):321–354, 2000.
- [4] U. G. Abdulla. On the Dirichlet problem for reaction-diffusion equations in nonsmooth domains. *Proceedings of the Third World Congress of Nonlinear Analysts, Part 2 (Catania, 2000)*, 47:765–776, 2001.
- [5] U. G. Abdulla. On the Dirichlet problem for the nonlinear diffusion equation in nonsmooth domains. *Journal of Mathematical Analysis and Applications*, 260(2):384–403, 2001.
- [6] U. G. Abdulla. Evolution of interfaces and explicit asymptotics at infinity for the fast diffusion equation with absorption. *Nonlinear Analysis*, 50(4):541–560, 2002.



- [7] U. G. Abdulla. Nonlinear diffusion in irregular domains. In *Elliptic and parabolic problems (Rolduc/Gaeta, 2001)*, pages 302–310. World Sci. Publ., River Edge, NJ, 2002. URL: [https://doi.org/10.1142/9789812777201\\_0029](https://doi.org/10.1142/9789812777201_0029).
- [8] U. G. Abdulla. First boundary value problem for the diffusion equation. I. Iterated logarithm test for the boundary regularity and solvability. *SIAM Journal on Mathematical Analysis*, 34(6):1422–1434, 2003. URL: <https://doi.org/10.1137/S0036141002415049>.
- [9] U. G. Abdulla. Kolmogorov problem for the heat equation and its probabilistic counterpart. *Nonlinear Analysis*, 63(5-7):712–724, 2005. URL: <https://doi.org/10.1016/j.na.2005.03.044>.
- [10] U. G. Abdulla. Multidimensional Kolmogorov-Petrovsky test for the boundary regularity and irregularity of solutions to the heat equation. *Boundary Value Problems*, (2):181–199, 2005.
- [11] U. G. Abdulla. Well-posedness of the Dirichlet problem for the nonlinear diffusion equation in non-smooth domains. *Transactions of the American Mathematical Society*, 357(1):247–265, 2005.
- [12] U. G. Abdulla. Necessary and sufficient condition for uniqueness of solution to the first boundary value problem for the diffusion equation in unbounded domains. *Nonlinear Analysis*, 64(5):1012–1017, 2006. URL: <https://doi.org/10.1016/j.na.2005.04.055>.
- [13] U. G. Abdulla. Reaction-diffusion in nonsmooth and closed domains. *Boundary Value Problems*, 2007(1):031261, 2006.

- [14] U. G. Abdulla. Wiener's criterion for the unique solvability of the Dirichlet problem in arbitrary open sets with non-compact boundaries. *Nonlinear Analysis*, 67(2):563–578, 2007.
- [15] U. G. Abdulla. Wiener's criterion at  $\infty$  for the heat equation. *Advances in Differential Equations*, 13(5-6):457–488, 2008.
- [16] U. G. Abdulla. Wiener's criterion at  $\infty$  for the heat equation and its measure-theoretical counterpart. *Electronic Research Announcements in Mathematical Sciences*, 15:44–51, 2008.
- [17] U. G. Abdulla. Regularity of  $\infty$  for the heat equation and the well-posedness of the Dirichlet problem. In *Advances in nonlinear analysis: theory methods and applications*, volume 3 of *Math. Probl. Eng. Aerosp. Sci.*, pages 173–180. Camb. Sci. Publ., Cambridge, 2009.
- [18] U. G. Abdulla. On the optimal control of the free boundary problems for the second order parabolic equations. I. Well-posedness and convergence of the method of lines. *Inverse Problems and Imaging*, 7(2):307–340, 2013.
- [19] U. G. Abdulla. On the optimal control of the free boundary problems for the second order parabolic equations. II. Convergence of the method of finite differences. *Inverse Problems and Imaging*, 10(4):869–898, 2016.
- [20] U. G. Abdulla, J. Du, A. Prinkey, C. Ondracek, and S. Parimoo. Evolution of interfaces for the nonlinear double degenerate parabolic equation of turbulent filtration with absorption. *Mathematics and Computers in Simulation*, 153:59–82, 2018.
- [21] U. G. Abdulla and R. Jeli. Evolution of interfaces for the non-linear parabolic  $p$ -Laplacian type diffusion equation of non-Newtonian elastic filtration with strong

- absorption. *European Journal of Applied Mathematics*, March 2019. URL: <https://doi.org/10.1017/S095679251900007X>.
- [22] U. G. Abdulla and R. Jeli. Evolution of interfaces for the non-linear parabolic  $p$ -Laplacian type reaction-diffusion equations. *European Journal of Applied Mathematics*, 28(5), 2017.
- [23] U. G. Abdulla and J. R. King. Interface development and local solutions to reaction-diffusion equations. *SIAM Journal on Mathematical Analysis*, 32(2):235–260, 2000.
- [24] U. G. Abdulla, A. Prinkey, and M. Avery. Evolution of interfaces for the nonlinear double degenerate parabolic equation of turbulent filtration with absorption. II. Fast diffusion case. *Submitted*, 2019. URL: <http://arxiv.org/abs/1903.08155>.
- [25] U. G. Abdullaev. Unbounded solutions of a nonlinear heat equation with a sink. *Zh. Vychisl. Mat. i Mat. Fiz.*, 32(8):1244–1257, 1992.
- [26] U. G. Abdullaev. On existence of unbounded solutions of nonlinear heat equations with absorption. *Zh. Vychisl. Mat. i Mat. Fiz.*, 33:232–245, 1993.
- [27] U. G. Abdullaev. On the localization of unbounded solutions of the nonlinear heat equation with transfer. *Dokl. Akad. Nauk*, 329(5):535–537, 1993.
- [28] U. G. Abdullaev. Large-time behaviour of solutions of the nonlinear infiltration equation. *Nonlinear Analysis*, 23(10):1353–1364, 1994. URL: [https://doi.org/10.1016/0362-546X\(94\)90153-8](https://doi.org/10.1016/0362-546X(94)90153-8).

- [29] U. G. Abdullaev. The space localization of unbounded boundary perturbations in nonlinear heat conduction with transfer. *Applied Mathematics Letters*, 7(6):91–95, 1994. URL: [https://doi.org/10.1016/0893-9659\(94\)90100-7](https://doi.org/10.1016/0893-9659(94)90100-7).
- [30] U. G. Abdullaev. Exact local estimates for the supports of solutions in problems for non-linear parabolic equations. *Russian Academy of Sciences Sbornik Mathematics*, 186(8), 1995.
- [31] U. G. Abdullaev. On asymptotically sharp local estimates for finite solutions of a nonlinear parabolic equation with absorption. *Sibirsk. Mat. Zh.*, 36(5):975–991, i, 1995. URL: <https://doi.org/10.1007/BF02112527>.
- [32] U. G. Abdullaev. On sharp local estimates for the support of solutions in problems for nonlinear parabolic equations. *Mat. Sb.*, 186(8):3–24, 1995. URL: <https://doi.org/10.1070/SM1995v186n08ABEH000058>.
- [33] U. G. Abdullaev. Local structure of solutions of the reaction-diffusion equations. *Nonlinear Analysis*, 30:3153–3163, 1997.
- [34] U. G. Abdullaev. Instantaneous shrinking and exact local estimations of solutions in nonlinear diffusion absorption. *Advances in Mathematical Sciences and Applications*, 8(1):483–500, 1998.
- [35] U. G. Abdullaev. Instantaneous shrinking of the support of a solution of a nonlinear degenerate parabolic equation. *Mat. Zametki*, 63(3):323–331, 1998. URL: <https://doi.org/10.1007/BF02317772>.
- [36] M. Agueh, A. Blanchet, and J. A. Carrillo. Large time asymptotics of the doubly nonlinear equation in the non-displacement convexity regime. *Journal of Evolution Equations*, 10(1):59–84, 2009.

- [37] G. Akagi and U. Stefanelli. A variational principle for doubly nonlinear evolution. *Applied Mathematics Letters*, 23(9):1120–1124, 2010.
- [38] S. Antontsev and S. Shmarev. Doubly degenerate parabolic equations with variable nonlinearity I: Existence of bounded strong solutions. *Advances in Differential Equations*, 17(11/12):1181–1212, 2012.
- [39] S. Antontsev and S. Shmarev. Parabolic equations with double variable nonlinearities. *Mathematics and Computers in Simulation*, 81(10):2018–2032, 2011.
- [40] S. N. Antontsev, J. I. Diaz, and S. Shmarev. *Energy Methods for Free Boundary Problems: Applications to Nonlinear PDEs and Fluid Mechanics*, volume 48. Springer Verlag, 2012.
- [41] M. Aripov and S. A. Sadullaeva. Qualitative properties of solutions of a doubly nonlinear reaction-diffusion system with a source. *Journal of Applied Mathematics and Physics*, 3:1090–1099, 2015.
- [42] G. I. Barenblatt. On some unsteady motions of a liquid or a gas in a porous medium. *Prikl. Mat. Mech.*, 16:67–78, 1952.
- [43] G. I Barenblatt. *Scaling, self-similarity, and intermediate asymptotics*. Cambridge Texts in Applied Mathematics. Cambridge University Press, 1996.
- [44] P. Benilan, M. G. Crandall, and M. Pierre. Solutions of the porous medium equation under optimal conditions on initial values. *Indiana University Mathematics Journal*, 33:51–87, 1984.
- [45] F. Bernis. Existence results for doubly nonlinear higher order parabolic equations on unbounded domains. *Mathematische Annalen*, 279(3):373–394, 1988.

- [46] T. M. Bokalo and O. M. Buhrii. Doubly nonlinear parabolic equations with variable exponents of nonlinearity. *Ukrainian Mathematical Journal*, 63(5):709–728, 2011.
- [47] C. Caisheng. Global existence and  $L^\infty$  estimates of solution for doubly nonlinear parabolic equation. *Journal of Mathematical Analysis and Applications*, 244(1):133–146, 2000.
- [48] P. Cianci, A. V. Martynenko, and A. F. Tedeev. The blow-up phenomenon for degenerate parabolic equations with variable coefficients and nonlinear source. *Nonlinear Analysis*, 7(1):2310–2323, 2010.
- [49] P. Colli. On some doubly nonlinear evolution equations in Banach spaces. *Japan Journal of Industrial and Applied Mathematics*, 9, 1992.
- [50] P. Colli and A. Visintin. On a class of doubly nonlinear evolution equations. *Communications in Partial Differential Equations*, 15(5):737–756, 1990.
- [51] J. I. Diaz and F. De Thelin. On a nonlinear parabolic problem arising in some models related to turbulent flows. *SIAM Journal on Mathematical Analysis*, 25(4):1085–1111, 1991.
- [52] E. DiBenedetto. *Degenerate Parabolic Equations*. Series Universitext. Springer Verlag, 1993.
- [53] E. DiBenedetto and M. A. Herrero. On the Cauchy problems and initial traces for a degenerate parabolic equation. *Transactions of the American Mathematical Society*, 314:187–224, 1989.

- [54] E. DiBenedetto and M. A. Herrero. Nonnegative solutions of the evolution  $p$ -Laplacian equations: Initial traces and Cauchy problem when  $1 < p < 2$ . *Archive for Rational Mechanics Analysis*, 111:225–290, 1990.
- [55] C. Ebmeyer and J. M. Urbano. Regularity in Sobolev spaces for doubly nonlinear parabolic equations. *Journal of Differential Equations*, 187(2):375–390, 2003.
- [56] C. Ebmeyer and J. M. Urbano. The smoothing property for a class of doubly nonlinear parabolic equations. *Transactions of the American Mathematical Society*, 357:3239–3253, 2005.
- [57] J. R. Esteban and J. L. Vazquez. On the equation of turbulent filtration in one-dimensional porous media. *Nonlinear Analysis*, 10(11):1303–1325, 1986.
- [58] S. Fornaro and M. Sosio. Intrinsic Harnack estimates for some doubly nonlinear degenerate parabolic equations. *Advances in Differential Equations*, 13(1-2):139–168, 2008.
- [59] L. Giacomelli and G. Grn. Lower bounds on waiting times for degenerate parabolic equations and systems. *Interfaces and Free Boundaries*, 8(1):111–129, 2006.
- [60] U. Gianazza and V. Vespri. A Harnack inequality for solutions of doubly nonlinear parabolic equations. *Journal of Applied Functional Analysis*, 2006.
- [61] B. H. Gilting and R. Kersner. A necessary and sufficient condition for finite speed of propagation in the theory of doubly nonlinear degenerate parabolic equations. *Proceedings of the Royal Society of Edinburgh Section A: Mathematics*, 126(4):739–767, 1996.

- [62] R. E. Grundy and L. A. Peletier. Short time behaviour of a singular solution to the heat equation with absorption. *Proc. Roy. Soc. Edinburgh Sect. A*, 107:271–288, 1987.
- [63] R. E. Grundy and L. A. Peletier. The initial interface development for a reaction-diffusion equation with power-law initial data. *The Quarterly Journal of Mechanics and Applied Mathematics*, 43(4):535–559, 1990.
- [64] M. A. Herrero and M. Pierre. The Cauchy problem for  $u_t = \delta u^m$  when  $0 < m < 1$ . *Transactions of the American Mathematical Society*, 291:145–158, 1985.
- [65] M. A. Herrero and J. L. Vazquez. Thermal waves in absorbing media. *Journal of Differential Equations*, 74:218–233, 1988.
- [66] K. Ishige. On the existence of solutions of the Cauchy problem for a doubly nonlinear parabolic equation. *SIAM Journal on Mathematical Analysis*, 27(5):1235–1260, 1996.
- [67] A. V. Ivanov. Hölder estimates for equations of slow and normal diffusion type. *Journal of Mathematical Sciences*, 85(1):1640–1644, 1997.
- [68] A. V. Ivanov. Regularity for doubly nonlinear parabolic equations. *Journal of Mathematical Sciences*, 83(1):22–37, 1997.
- [69] A. V. Ivanov, P. Z. Mkrtychan, and W. Jäger. Existence and uniqueness of a regular solution of the Cauchy-Dirichlet problem for a class of doubly nonlinear parabolic equations. *Journal of Mathematical Sciences*, 84(1):845–855, 1997.



- [70] W. Jäger and J. Kačur. Solution of doubly nonlinear and degenerate parabolic problems by relaxation schemes. *Mathematical Modeling and Numerical Analysis*, 29(5):605–627, 1995.
- [71] A. S. Kalashnikov. The influence of absorption on the propagation of heat in a medium with heat conductivity that depends on the temperature. *Zh. Vychisl. Mat. i Mat. Fiz.*, 16:689–696, 1976.
- [72] A. S. Kalashnikov. On a nonlinear equation appearing in the theory of non-stationary filtration. *Trud. Semin. I. G. Pertovski*, 4:137–146, 1978.
- [73] A. S. Kalashnikov. On the propagation of perturbations in the first boundary value problem of a doubly-nonlinear degenerate parabolic equation. *Trud. Semin. I. G. Pertovski*, 8:128–134, 1982.
- [74] A. S. Kalashnikov. Some problems of the qualitative theory of non-linear degenerate second-order parabolic equations. *Russian Mathematical Surveys*, 42(2):169–222, 1987.
- [75] S. Kamin, L. A. Peletier, and J. L. Vazquez. A nonlinear diffusion-absorption equation with unbounded initial data. pages 243–263, 1992.
- [76] J. Kinnunen and T. Kuusi. Local behaviour of solutions to doubly nonlinear parabolic equations. *Mathematische Annalen*, 337(3):705–728, 2007.
- [77] I. Kombe. Doubly nonlinear parabolic equations with singular lower order term. *Nonlinear Analysis*, 56(2):185–199, 2004.
- [78] L. S. Leibenson. General problem of the movement of a compressible fluid in porous medium. *Izv. Akad. Nauk SSSR, Geography and Geophysics*, IX:7–10, 1945.

- [79] J. Li. Cauchy problem and initial trace for a doubly degenerate parabolic equation with strongly nonlinear sources. *Journal of Mathematical Analysis and Applications*, 264:49–67, 2001.
- [80] Z. Li, W. Du, and C. Mu. Travelling-wave solutions and interfaces for non-Newtonian diffusion equations with strong absorption. *Journal of Mathematical Research with Applications*, 334:451–462, 2013.
- [81] Y. G. Lu and L. Qian. Regularity of viscosity solutions of a degenerate parabolic equation. *Proceedings of the American Mathematical Society*, 2002.
- [82] J. J. Manfredi and V. Vespri. Large time behavior of solutions to a class of doubly nonlinear parabolic equations. *Electronic Journal of Differential Equations*, 1994(02):1–17, 1994.
- [83] A. Matas and J. Merker. Existence of weak solutions to doubly degenerate diffusion equations. *Applications of Mathematics*, 57(1):43–69, 2012.
- [84] C. Mu, P. Zheng, and D. Liu. Localization of solutions to a doubly degenerate parabolic equation with a strongly nonlinear source. *Communications in Contemporary Mathematics*, 14(03), 2012.
- [85] O. A. Oleinik, A. S. Kalashnikov, and C. Y. Lin. Cauchy problem and boundary value problems for an equation of nonstationary filtration. *Izv. Akad. Nauk SSSR, Ser. Mat.*, 22:667–704, 1958.
- [86] M. Ôtani and Y. Sugiyama. Lipschitz continuous solutions of some doubly nonlinear parabolic equations. *Discrete and Continuous Dynamical Systems*, 8(3):647–670, 2002.

- [87] S. Shmarev, V. Vdovin, and A. Vlasov. Interfaces in diffusion-absorption processes in nonhomogeneous media. *Mathematics and Computers in Simulation*, 2015.
- [88] S. N. Antontsev and S. I. Shmarev. Doubly degenerate parabolic equations with variable nonlinearity II: Blow-up and extinction in a finite time. *Nonlinear Analysis*, 95:483–498, 2014.
- [89] M. Slodicka. A robust and efficient linearization scheme for doubly nonlinear and degenerate parabolic problems arising in flow in porous media. *SIAM Journal on Scientific Computing*, 23(5):1593–1614, 2012.
- [90] U. Stefanelli. On a class of doubly nonlinear nonlocal evolution equations. *Differential and Integral Equations*, 15(8):897–922, 2002.
- [91] A. F. Tedeev. The interface blow-up phenomenon and local estimates for doubly degenerate parabolic equations. *Applicable Analysis*, 86(6):755–782, 2007.
- [92] M. Tsutsumi. On solutions of some doubly nonlinear degenerate parabolic equations with absorption. *Journal of Mathematical Analysis and Applications*, 132(1):187–212, 1988.
- [93] C. J. van Duijn and Z. Hongfei. Regularity properties of a doubly degenerate equation in hydrology. *Communications in Partial Differential Equations*, 13(3):261–319, 1988.
- [94] J. L. Vazquez. *The Porous Medium Equation: Mathematical Theory*. Oxford Science Publications. Oxford University Press, 2007.
- [95] V. Vespri. On the local behaviour of solutions of a certain class of doubly nonlinear parabolic equations. *Manuscripta Mathematica*, 75(1):65–80, 1992.

- [96] G. L. Yun. Hölder estimates of solutions to some doubly nonlinear degenerate parabolic equations. *Communications in Partial Differential Equations*, 24(5-6):895–913, 1999.
- [97] Ya. B. Zeldovich and A. S. Kompaneets. On the theory of heat propagation for temperature dependent thermal conductivity, in collection commemorating the 70th anniv. of A. F. Ioffe. *Izdat. Akad. Nauk SSSR*, 1950.
- [98] H. Zhan. Large time behavior of solutions to a class of doubly nonlinear parabolic equations. *Applications of Mathematics*, 53(6):521–533, 2008.
- [99] H. Zhan. The asymptotic behavior of solutions for a class of doubly degenerate nonlinear parabolic equations. *Journal of Mathematical Analysis and Applications*, 370(1):1–10, 2010.
- [100] H. Zhan. The asymptotic behavior of a doubly nonlinear parabolic equation with a absorption term related to the gradient. *WSEAS Transactions on Mathematics*, 2011.
- [101] J. Zhao and H. Yuan. The Cauchy problem of some doubly nonlinear degenerate parabolic equations. *Chinese Science Abstracts*, 1995.
- [102] T. Zheng and J. Zhao. On the Cauchy problem for a doubly nonlinear degenerate parabolic equation with strongly nonlinear sources. *Science in China Series A: Mathematics*, 51(11):2059–2071, 2008.

# Chapter 6

## Appendix

### 6.1

Here we give explicit values of the constants used in Chapter 2 ( $mp > 1$ ).

I.  $0 < \beta < 1$  and  $\alpha < (1+p)/(mp - \min\{1, \beta\})$ .

$$\xi_1 = p^{\frac{1}{1+p}} (\alpha(mp-1))^{-\frac{1}{1+p}}, \xi_2 = 1, p(mp-1)^{-1} \leq \alpha < (1+p)(mp-1)^{-1},$$

$$\xi_1 = 1, \xi_2 = p^{\frac{1}{1+p}} (\alpha(mp-1))^{-\frac{1}{1+p}}, 0 < \alpha \leq p(mp-1)^{-1}.$$

II.  $0 < \beta < 1$  and  $\alpha = (1+p)/(mp - \beta)$ .

$$f_1(0) = A_1 > 0,$$

$$\zeta_1 = A_1^{\frac{mp-1}{1+p}} (1 + b(1-\beta)A_1^{\beta-1})^{-\frac{1}{1+p}} (p(mp)^p(1-\beta))^{\frac{1}{1+p}} (mp-1)^{-1},$$

$$C_1 = A_1 \zeta_1^{-\frac{p}{mp-1}}, p(m+\beta) > 1+p, C > C_*,$$

$$\zeta_1 = A_1^{\frac{mp-1}{1+p}} (1+b(1-\beta)A_1^{\beta-1})^{-\frac{1}{1+p}} ((m(1+p))^p p(m+\beta)(1-\beta))^{\frac{1}{1+p}} (mp-\beta)^{-1},$$

$$C_1 = A_1 \zeta_1^{-\frac{1+p}{mp-\beta}}, p(m+\beta) < 1+p, C > C_*,$$

$$\zeta_2 = A_1^{\frac{mp-1}{1+p}} (1+b(1-\beta)A_1^{\beta-1})^{-\frac{1}{1+p}} ((m(1+p))^p p(m+\beta)(1-\beta))^{\frac{1}{1+p}} (mp-\beta)^{-1},$$

$$C_2 = A_1 \zeta_2^{-\frac{1+p}{mp-\beta}}, p(m+\beta) > 1+p, C > C_*,$$

$$\zeta_2 = (A_1/C_*)^{\frac{mp-\beta}{1+p}}, C_2 = C_*, p(m+\beta) < 1+p, C > C_*,$$

$$\zeta_1 = -C^{-\frac{mp-\beta}{1+p}} (b(1-\beta))^{\frac{mp-\beta}{(1+p)(1-\beta)}}, p(m+\beta) > 1+p, 0 < C < C_*,$$

$$\zeta_2 = -C^{-\frac{mp-\beta}{1+p}} \left( b(1-\beta)(1-(C/C_*)^{mp-\beta})^{\frac{mp-\beta}{(1+p)(1-\beta)}} \right), p(m+\beta) < 1+p, 0 < C < C_*,$$

$$R_1 = (m(1+p))^p p(1+p-p(m+\beta))(b(mp-\beta))^{1+p}^{-1},$$

$$R_2 = (m(1+p))^p (1+p)p(m+\beta-1)(b(mp-\beta))^{1+p}^{-1},$$

$$\theta_* = \left[ 1 - (C/C_*)^{mp-\beta} \right] \left[ (C_*/C)^{\frac{(mp-\beta)(1-\beta)}{1+p-p(m+\beta)}} - 1 \right]^{-1},$$

$$\ell_0 = C_*^{-\frac{mp-\beta}{1+p}} (C_*/C)^{\frac{(mp-\beta)(1-\beta)}{1+p-p(m+\beta)}} (b(1-\beta)\theta_*)^{\frac{mp-\beta}{(1+p)(1-\beta)}},$$

$$\zeta_3 = C_*^{-\frac{mp-\beta}{1+p}} \left[ (C_*/C)^{\frac{(mp-\beta)(1-\beta)}{1+p-p(m+\beta)}} - 1 \right] (b(1-\beta)\theta_*)^{\frac{mp-\beta}{(1+p)(1-\beta)}},$$

$$\ell_1 = C^{-\frac{mp-\beta}{1+p}} \left[ b(1-\beta)(\delta_*\Gamma)^{-1} (1-\delta_*\Gamma - (1-\delta_*\Gamma)^{-p}(C/C_*)^{mp-\beta}) \right]^{\frac{mp-\beta}{(1+p)(1-\beta)}},$$

$$\zeta_4 = \delta_*\Gamma\ell_1, \Gamma = 1 - (C/C_*)^{\frac{mp-\beta}{1+p}}, C_3 = C(1-\delta_*\Gamma)^{-\frac{1+p}{mp-\beta}},$$

where  $\delta_* \in (0, 1)$  satisfies

$$g(\delta_*) = \max_{[0,1]} g(\delta), g(\delta) = \delta^{\frac{1+p-p(m+\beta)}{mp-\beta}} (1-\delta\Gamma - (1-\delta\Gamma)^{-p}(C/C_*)^{mp-\beta}).$$

III.  $0 < \beta < 1$  and  $\alpha > (1 + p)/(mp - \beta)$ .

$$\zeta_5 = (\ell_*/\ell)^{\alpha(1-\beta)}(1 - \varepsilon)\ell,$$

$$C_6 = [1 - (\ell_*/\ell)^{\alpha(1-\beta)}(1 - \varepsilon)]^{-\alpha} [C^{1-\beta} - \ell^{-\alpha(1-\beta)}b(1 - \beta)(1 - \varepsilon)]^{1/(1-\beta)}.$$

V.  $b = 0$  and  $\alpha > 0$ .

$$w(0, 1) = A_0 > 0,$$

$$\bar{C} = \left[ \frac{(mp - 1)^{1+p}}{p(m+1)(m(1+p))^p} \right]^{\frac{1}{mp-1}},$$

$$\gamma_\varepsilon = \frac{p(m+1)(m(1+p))^p(C + \varepsilon)^{mp-1}}{(mp - 1)^p} + \varepsilon,$$

$$\xi_3 = A_0^{\frac{mp-1}{1+p}} \left[ \frac{(mp)^p(1+p - \alpha(mp-1))}{(mp-1)^p} \right]^{\frac{1}{1+p}} C^{\frac{mp-1}{1+p - \alpha(mp-1)}} \xi_1,$$

$$\xi_4 = A_0^{\frac{mp-1}{1+p}} \left[ \frac{(mp)^p(1+p - \alpha(mp-1))}{(mp-1)^p} \right]^{\frac{1}{1+p}} C^{\frac{mp-1}{1+p - \alpha(mp-1)}} \xi_2,$$

$$C_4 = C^{(1+p)/(1+p - \alpha(mp-1))} A_0 \xi_3^{p/(1-mp)}, \quad C_5 = C^{(1+p)/(1+p - \alpha(mp-1))} A_0 \xi_4^{p/(1-mp)}.$$

## 6.2

Here we give explicit values of the constants used in Chapter 4 ( $0 < mp < 1$ ).

I.  $0 < \beta < mp$  and  $0 < \alpha < (1 + p)/(mp - \beta)$ .

$$z_1 = (b(1 - \beta))^{\frac{mp-1}{(1+p)(1-\beta)}} (m(1+p))^{\frac{p}{1+p}} (p(m+\beta))^{\frac{1}{1+p}} (mp - \beta)^{\frac{p(m+\beta)-1}{(1+p)(1-\beta)}} (1 - mp)^{\frac{1-mp}{(1+p)(1-\beta)}},$$

$$C_1 = \left( \frac{1 - \beta}{1 - mp} \right)^{\frac{1}{mp-\beta}} C_*,$$

$$\tau_0 = \left( \frac{1-mp}{mp-\beta} \right)^{\frac{mp-1}{1-\beta}} \left( \frac{\mathcal{D}}{C_*} \right)^{\frac{(mp-1)(\beta-mp)}{(1+p)(1-\beta)}}, \quad z_2 = \tau_0 \frac{1-\beta}{mp-\beta}.$$

II.  $0 < \beta < mp$  and  $\alpha = (1+p)/(mp-\beta)$ .

$$C_2 = A_1 z_3^{\frac{1+p}{\beta-mp}}, \quad A_1 = f_1(0) > 0,$$

$$z_3 = (m(1+p))^{\frac{p}{p+1}} (m+\beta)^{\frac{1}{1+p}} p^{\frac{1}{1+p}} (mp-\beta)^{-1} (1-\beta)^{\frac{1}{1+p}} A_1^{\frac{m-1}{1+p}} \left[ b(1-\beta) A_1^{\beta-1} + 1 \right]^{\frac{-1}{1+p}},$$

$$z_4 = \left( \frac{A_1}{C_*} \right)^{\frac{mp-\beta}{1+p}}, \quad z_5 = \tau_1 - \left( \frac{\rho}{C_*} \right)^{\frac{1+p}{mp-\beta}}, \quad C_3 = C \left( \frac{1}{1-\delta_* \Gamma} \right)^{\frac{1+p}{mp-\beta}},$$

$$\Gamma = 1 - \left( \frac{C}{C_*} \right)^{\frac{mp-\beta}{1+p}}, \quad z_6 = \delta_* \Gamma \tau_2, \quad \text{with } \delta_* \text{ such that } g(\delta_*) = \max_{\delta \in (0,1)} g(\delta),$$

$$g(\delta) = (\delta \Gamma)^{\frac{1+p-p(m+\beta)}{(1+p)(1-\beta)}} \left[ 1 - \delta \Gamma - \left( \frac{C}{C_*} \right)^{mp-\beta} \left( \frac{1}{1-\delta \Gamma} \right)^p \right]^{\frac{mp-\beta}{(1+p)(1-\beta)}},$$

$$\tau_2 = C^{\frac{\beta-mp}{1+p}} \left[ \frac{b(1-\beta)}{\delta_* \Gamma} \left( 1 - \delta_* \Gamma - \left( \frac{C}{C_*} \right)^{mp-\beta} \left( \frac{1}{1-\delta_* \Gamma} \right)^p \right) \right]^{\frac{mp-\beta}{(1+p)(1-\beta)}}.$$

V.  $\beta > mp$ .

$$C_5 = (1-\varepsilon)^{\frac{1}{1-mp}} \mathcal{D},$$

$$C_6 = \left( \frac{\alpha(1-mp)^{p+1}}{\varkappa_b(1+p+\alpha(1-mp))(m(1+p))^p p(m+1)} \right)^{\frac{1}{mp-1}},$$

$$\chi_1 = (A_0 - \varepsilon)^{(mp-1)/(1+p)} (1-\varepsilon)^{1/(1+p)} \mathcal{D}^{(1-mp)/(1+p)}, \quad \text{if } b > 0, 1 \leq \beta < (p(1-m)+2)/(1+p),$$

$$\chi_1 = (A_0 - \varepsilon)^{(mp-1)/(1+p)} \mathcal{D}^{(1-mp)/(1+p)}, \quad \text{if } b > 0, \beta \geq (p(1-m)+2)/(1+p) \text{ or } b < 0, \beta \geq 1,$$

$$\chi_2 = \left( \frac{A_0 + \varepsilon}{C_6} \right)^{\frac{mp-1}{1+p}}, \quad A_0 = f(0) > 0, \quad \varkappa_b = \begin{cases} 1, & \text{if } b > 0, \\ 1 + \varepsilon, & \text{if } b < 0, \end{cases}$$



$$z_8 = \left[ b(1-\beta)C_*^{\beta-1}(1-\varepsilon)^{mp-1}(1-(1-\varepsilon)^{\beta-mp}) \right]^{\frac{mp-\beta}{(1+p)(1-\beta)}},$$

$$\chi_3 = (A_0/\mathcal{D})^{(mp-1)/(1+p)},$$

$$\chi_4 = \chi_3(1+(1+p)/\alpha(1-mp))^{1/(1+p)},$$

$$C_7 = \mathcal{D}(1+(1+p)/\alpha(1-mp))^{1/(1+mp)}.$$